



HHS Public Access

Author manuscript

Cell Stem Cell. Author manuscript; available in PMC 2019 July 05.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Published in final edited form as:

Cell Stem Cell. 2018 July 05; 23(1): 25–30. doi:10.1016/j.stem.2018.04.004.

Human adult neurogenesis: evidence and remaining questions

Gerd Kempermann¹, Fred H. Gage², Ludwig Aigner³, Hongjun Song⁴, Maurice Curtis⁵, Sandrine Thuret⁶, H. Georg Kuhn⁷, Sebastian Jessberger⁸, Paul W. Frankland⁹, Heather A. Cameron¹⁰, Elizabeth Gould¹¹, Rene Hen¹², D. Nora Abrous¹³, Nicolas Toni¹⁴, Alejandro F. Schinder¹⁵, Xinyu Zhao¹⁶, Paul J. Lucassen¹⁷, and Jonas Frisén¹⁸

¹German Center for Neurodegenerative Diseases (DZNE) Dresden, and CRTD – Center for Regenerative Therapies Dresden, Technische Universität Dresden, Germany ²Laboratory of Genetics, The Salk Institute for Biological Studies, La Jolla, USA ³Institute of Molecular Regenerative Medicine, Spinal Cord Injury and Tissue Regeneration Center Salzburg, Paracelsus Medical University, Salzburg, Austria ⁴Department of Neuroscience, University of Pennsylvania, Philadelphia, USA ⁵Department of Anatomy and Medical Imaging, and Centre for Brain Research, The University of Auckland, Auckland, New Zealand ⁶King's College London, Institute of Psychiatry, Psychology & Neuroscience, Department of Basic and Clinical Neuroscience, London, United Kingdom ⁷University of Gothenburg, Institute for Neuroscience and Physiology, Section for Clinical Neuroscience, Gothenburg, Sweden, and Charité – Universitätsmedizin Berlin, Neurocure Cluster of Excellence, Berlin, Germany ⁸HiFo/Brain Research Institute, University of Zurich, Zurich, Switzerland ⁹Program in Neuroscience and Mental Health, Hospital for Sick Children, Toronto, ON M5G0A4, Canada ¹⁰Section on Neuroplasticity, National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20892, USA ¹¹Princeton Neuroscience Institute, Princeton University, Princeton, New Jersey ¹²Departments of Neuroscience and Psychiatry, Columbia University, New York, NY ¹³Neurocentre Magendie, INSERM U1215, Bordeaux, France ¹⁴Lausanne University Hospital, Department of Psychiatry, Center for Psychiatric Neurosciences, Lausanne, Switzerland ¹⁵Neuronal Plasticity Lab, Leloir Institute – CONICET, Buenos Aires, Argentina ¹⁶Waisman Center and Department of Neuroscience, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA ¹⁷Brain Plasticity group, Swammerdam Institute for Life Sciences, Center for Neuroscience, University of Amsterdam, The Netherlands ¹⁸Department of Cell and Molecular Biology, Karolinska Institute, Stockholm, Sweden

Abstract

Renewed discussion about whether or not adult neurogenesis exists in the human hippocampus, and the nature and strength of the supporting evidence, has been reignited by two prominently published reports with opposite conclusions. Here, we summarize the state of the field and argue

Correspondence should be directed to: gerd.kempermann@dzne.de (lead author), jonas.frisén@ki.se or gage@salk.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest statement: The authors declare no financial or non-financial conflict of interest.

that there is currently no reason to abandon the idea that adult-generated neurons make important functional contributions to neural plasticity and cognition across the human life span.

Graphical abstract

Kempermann et al. summarize the state of the adult neurogenesis field and argue that there is currently no reason to abandon the idea that adult-generated neurons make important functional contributions to neural plasticity and cognition across the human life span

Adult hippocampal neurogenesis, the lifelong generation of new neurons in a brain region that is central to learning and memory (Altman and Das, 1965), exerts a strong fascination for scientists and the public alike. Knowledge about this process has fundamentally changed our ideas about how the hippocampus works and, by extension, about the structural substrates that underlie human cognition, cognitive aging, and the loss of hippocampal functions in, for example, Alzheimer's disease or stress-related disorders and depression.

Two prominently published studies have now re-ignited the scientific debate about adult neurogenesis in humans. A report by Sorrells et al. (2018) concluded that neurogenesis in the human hippocampal dentate gyrus drops to negligible amounts during childhood, and that the human hippocampus must function differently from that in other species, in which adult neurogenesis is conserved (Sorrells et al., 2018). In another study, Boldrini et al. (2018) came to the opposite conclusion and reported lifelong neurogenesis in humans (Boldrini et al., 2018). Thus, in the space of only a few weeks, two reports have been published that could not be more different. Herein, we discuss how the current state of knowledge about adult hippocampal neurogenesis applies to the human situation (Fig. 1).

The evidence for adult neurogenesis in the human brain

In 1998, Eriksson and colleagues applied the current 'gold standard' adult hippocampal neurogenesis method, which was previously established in animal studies, on the human hippocampus (Eriksson et al., 1998). They identified patients who had received infusions of the thymidine analog, bromodeoxyuridine (BrdU) for tumor staging purposes, but did not receive any treatment that is thought to affect cell generation, and they analyzed the brains post mortem. Their conclusion from five brains was that adult neurogenesis could be detected in the human hippocampus in the same location and numbers as expected based on work in rats. BrdU and other halogenated thymidine analogs, such as IdU or CldU, are incorporated into the DNA of dividing precursor cells and can be detected immunohistochemically. Detecting a BrdU-positive neuron thus indicates that that neuron has originated from a cell that underwent division at exactly the time at which BrdU was applied, since BrdU has a short biological half-life. BrdU is not significantly incorporated during DNA repair and is not taken up by dying neurons (Bauer and Patterson, 2005). While such birthdating methods are cornerstones of demonstrating adult neurogenesis, especially in undescribed regions of the brain or in new species, they alone are not sufficient as proof but require support by methodologically independent lines of evidence.

Author Manuscript
Author Manuscript
Author Manuscript
Author Manuscript

Providing such supporting evidence, stem cells with neurogenic potential were isolated from the adult human hippocampus (e.g., (Palmer et al., 2001)). In addition, several studies have used immunocytochemistry to detect cells expressing cell proliferation markers in human postmortem brains (e.g., (Boekhoorn et al., 2006; Curtis et al., 2003; Dennis et al., 2016; Liu et al., 2008; Mathews et al., 2017)).

Both Sorrells et al. and Boldrini et al. primarily base their main conclusions on the individual or combined expression of key marker proteins such as doublecortin (DCX) or PSA-NCAM as markers for intermediate progenitor cells and early immature neurons (often dubbed ‘neuroblasts’). In rodents, DCX (and PSA-NCAM) characterizes an intermediate phase of adult neurogenesis between the precursor cell stage and immature neurons, and it is widely used as a proxy marker for adult neurogenesis, although it is also expressed in other contexts (Kuhn et al., 2016). Several earlier studies have used DCX to assess adult neurogenesis in humans (Dennis et al., 2016; Galán et al., 2017; Knoth et al., 2010; Liu et al., 2008; Mathews et al., 2017).

Sorrells et al. and Boldrini et al. use the approach first reported by Knoth et al. (2010), who assessed 54 samples across the life span of 0 to 100 years using combinations of 14 markers (Knoth et al., 2010). In contrast to Sorrells et al. (2018), Knoth et al. and now Boldrini et al. found DCX-positive cells co-expressing other neurogenesis markers. But while Sorrells et al. and several other studies pointed out an age-related decrease in marker overlap and a sharp decline in proliferating cells (Dennis et al., 2016; Knoth et al., 2010; Mathews et al., 2017), Boldrini et al. state that these numbers would hardly change across adulthood. In contrast to previous studies, they applied stereology, a method for unbiased quantification within a tissue volume, so their argument carries more weight. The conclusion still stands in contrast to quantitative estimates, based on carbon 14 (^{14}C) birthdating of neuronal DNA (Spalding et al., 2013). That study by Spalding, Bergmann, and colleagues, assessing ^{14}C data from 55 individuals, is widely seen as the long-awaited independent proof that adult neurogenesis does indeed exist in the human hippocampus. Another study by the same group, though focused on striatal neurogenesis, also contained a replication of Eriksson’s findings using the thymidine analog IdU in four more subjects (Fig. S2 of (Ernst et al., 2014)).

The studies by Eriksson et al., Ernst et al. and Spalding et al. used a form of lineage tracing in which the DNA of dividing precursor cells was labeled (either by ^{14}C , BrdU, or IdU) and their progeny was analyzed for the expression of neuronal markers. While histological marker studies, like Sorrells et al., based their conclusion about neurogenesis on the starting point of adult neurogenesis, or the precursor cells, their proliferation, and early immature stages, these other studies focused on the end product, or the actual demonstration of the presence of newly formed neurons.

Technical issues

The limitations of marker studies

Sorrells et al. essentially based their conclusion about the absence of neurogenesis on the absence of morphological features and the lack of detecting two marker proteins, DCX and

PSA-NCAM. Boldrini et al. used the same markers as evidence of neurogenesis. A crucial factor in accurate detection of marker proteins is the postmortem delay (PMD), i.e., the time between the death of a person and fixation of the brain. DCX rapidly breaks down after death: a controlled time course study of PMD in rats has shown that DCX staining becomes weak within a few hours of PMD (Boekhoorn et al., 2006). In the Sorrells paper, many subjects had very long PMDs of ‘less than 48 hrs’ (that is, up to 2 full days before fixation), whereas in Boldrini et al. (2018) the PMD was up to 26 hours. It would be valuable to know whether the inter-individual variation in neuroblast numbers in Boldrini et al. correlated to the PMD. In addition, it would be interesting to learn how well the different marker combinations (and, hence, cell types) correlated. Type and duration of fixation are other relevant methodological factors. Human samples might be stored in the fixative for years. Based on their long fixation period in 10% formalin,, masking of the PSA-NCAM antigen has likely occurred, possibly explaining the relative absence of this marker in in Sorrells et al.’s tissues.

The disease phase preceding the study subjects’ deaths also needs to be considered. Moreover, in humans, the act of dying itself massively elevates stress hormones (Bao and Swaab, 2018) and since DCX staining dropped dramatically as soon as 30 minutes after capture in bats, stress hormones may have reduced DCX levels in the human brain as well (Chawana et al., 2014).

Variability can also be explained by the many genetic and environmental factors that regulate neurogenesis in rodents such as exercise, hormonal status, diet, epilepsy, anxiety, addiction, inflammation, and stress (Lucassen et al., 2015). The study by Boldrini et al. differs from previous studies in that it attempts to correlate adult neurogenesis with angiogenesis and tissue volume as additional tissue parameters, which are influenced by ‘activity.’ Moreover, they included only subjects without neurological or psychiatric disease.

Given these methodological issues and the impact of ‘lifestyle’ factors for the human tissues that were studied at an ‘end stage,’ it seems likely that the Sorrells et al.’s study was at least not optimized for the detection of neurogenesis.

An important consequence of the renewed discussion will therefore be a raised awareness of the challenges that these approaches pose when studying human brains (Bao and Swaab, 2018).

Quantitative aspects

Several groups have previously reported quantitative estimates of the presence of DCX- or PSA-NCAM-positive cells in adult humans (Dennis et al., 2016; Galán et al., 2017; Knoth et al., 2010), and Boldrini and colleagues (2018) have been among the first to make a serious attempt to apply proper stereological principles to the analysis. This approach is urgently needed, but the implementation is challenging in the kind of tissue samples usually available from humans. Irrespective of the approach used, all of these studies reported only sparse DCX-positive cells in the adult dentate gyrus, and the rough quantitative estimates actually seem comparable between the studies.

Carbon dating indicates that about 700 new neurons are added per day in each dentate gyrus and it seems that, even if one allows a large margin of error, the available numbers for DCX-expressing cells fall into the same order of magnitude. The decline in the number of DCX-positive cells during adulthood and into old age, reported in most studies, is closely paralleled by a decreased generation of new neurons measured by carbon dating (Fig 5A in (Spalding et al., 2013). This decline is also found in rodents, where not only proliferation decreases but also the subsequent neurogenesis phase slows down with increasing age. If the numbers from Boldrini et al. are confirmed, the extent of adult human neurogenesis would previously have been under-estimated rather than over-estimated.

For the number of DCX-positive cells found by Knoth et al. to give rise to the number of new neurons estimated by carbon dating, the phase of DCX expression could last for approximately three weeks if half of them gave rise to mature neurons. This duration of the DCX-positive stage is comparable to what is seen in rodents, in which approximately half of the DCX-positive intermediate cells give rise to a mature neuron. Thus, it is conceivable that the reported very sparse numbers of DCX-positive cells in the adult human dentate gyrus can still give rise to the number of new neurons quantified by the BrdU method and carbon dating. However, there is a large inter-individual variation in the number of ‘neuroblasts’ reported by Boldrini et al., with very low numbers in some subjects. Such inter-individual variation has been suggested by a previous marker study (Dennis et al., 2016), as well as by carbon dating (Spalding et al., 2013). It does not seem likely, but it is still conceivable that the individuals in the sample of the Sorrells et al. study all happened to have minimal or no neurogenesis.

Conceptual contexts

Potential species differences

The use of DCX and PSA-NCAM expression as sole indicators of ‘neurogenesis’ is also problematic as, in humans, we might find a relative temporal ‘decoupling’ of precursor cell proliferation, which builds the potential for neurogenesis, from the actual recruitment or differentiation into new neurons. One study suggested, for example, that the decrease in DCX in the aging human hippocampus is not paralleled by similar decreases in proliferation marker Ki67, putative stem cell marker GFAP8, or neurogenic transcription factor Tbr2/EOMES (Mathews et al., 2017). The learning-induced recruitment of newborn neurons (at least in rodents) is dependent on a reservoir of recruitable postmitotic cells and not on precursor cell proliferation *per se*. DCX is often used as a proxy for this population of “immature” neurons. However, there is no simple relationship between cell proliferation, the number of DCX-positive cells, and net neurogenesis. In fact, DCX expression is not required for adult neurogenesis or synaptic plasticity during that period (Germain et al., 2013). DCX expression alone is thus likely not sufficient to fully understand the functional potential of neurogenesis.

In addition, new neurons in mice are not DCX-positive throughout their entire postmitotic maturation period, and rats have many fewer DCX-positive cells than mice, despite having higher rates of neurogenesis, because their neurons mature faster (Snyder et al., 2009). In mice, Calretinin (CR) appears to be a better proxy marker for this period. Ironically, CR

does not seem to be similarly expressed even in rats, but it has been used in at least one human study (Galán et al., 2017). It is clearly speculation at this time, but if DCX does not cover the entire period of increased plasticity in mice, we should be open to the possibility that species (as well as inter-individual) differences also apply to the dynamics of marker expression and the lengths of critical phases (Fig. 2).

Functional aspects

Research across many laboratories has painted an increasingly complete picture of how new neurons contribute to hippocampal function (Abrous and Wojtowicz, 2015; Christian et al., 2014). These studies support the view that adult neurogenesis is not needed for learning *per se* but rather for an advanced level of functionality. The new neurons allow the spatiotemporal contextualization of information and they help avoid catastrophic interference in the hippocampal network, promoting ‘behavioral pattern separation.’ They facilitate the integration of new information into pre-existing contexts and help to clear the dentate gyrus at the circuit level and, at least in this sense, support forgetting. In addition, as the hippocampus is part of the limbic system, new neurons are involved in affective behaviors.

The new neurons contribute synaptic plasticity to the dentate gyrus, measured as increased long-term potentiation (LTP; (Ge et al., 2007; Marín-Burgin et al., 2012; Schmidt-Hieber et al., 2004). All other neurons are massively inhibited by the local interneurons. At a given time, synaptic plasticity in the dentate gyrus is thus concentrated in a defined, functionally naive sub-set of (new) neurons. This unique mechanism of focusing plasticity sets this neuronal network apart from all others studied to date. In this context, the number of new cells required for a functional benefit is actually very low.

As an important and influential discussion point, Pasko Rakic has famously argued that adult hippocampal neurogenesis would not be possible in humans because the adult human brain has to favor stability over plasticity in order to accomplish its computational tasks (Rakic, 1985). Current theories usually argue the other way around: it is exactly its amazing plasticity that has made the human brain so flexible and successful. The question is: what is the contribution of new neurons to this success? As adult neurogenesis is spatially limited, this contribution cannot be general, as most brain regions, including the neocortex, work without it, but its effects might still be strategic. Simple brains are highly effective but in their “hard-wiredness,” they are hardly adaptable. Adult hippocampal neurogenesis is a prime tool for adaptability and seems to be a solution to the specific computational challenge in the dentate gyrus. Without it, yet another solution to the plasticity-stability dilemma as seen in rodents would need to have evolved in humans. Whether such a parallel solution is likely or not remains to be discussed but the functional contribution that new neurons would make to human cognition is not negligible.

Evolutionary considerations

The mammalian dentate gyrus as we see it in rodents and primates, including humans, is an “add-on” structure that evolved late phylogenetically and developed late ontogenetically. Signs of adult hippocampal neurogenesis have been detected across essentially all land-born

mammalian species, that is except for the aquatic and possibly some flying mammals (Kempermann, 2012). Dolphins, however, despite their ascribed ‘intelligence,’ have a habitat that is profoundly different from humans, and they have an exceptionally small hippocampus and a cortical architecture that differs massively from terrestrial mammals. By all standards, humans are more like mice in this respect.

Adult hippocampal neurogenesis evolved with the dentate gyrus; it shows little resemblance to the more diffuse neurogenesis found in the non-mammalian equivalents. Additional comparative studies are still needed, but the hypothesis is that adult hippocampal neurogenesis is an advanced solution to a particular network situation that delivers added specialized functionality to the hippocampus – including in humans. Sorrells et al. argue that there is no condensed subgranular zone stem cell niche or neurogenesis in humans and thus that such continuity in function might not exist, but this cannot be concluded from the presence or absence of marker proteins alone. The described functional relevance of adult neurogenesis is dependent on the availability of ‘immature’ neurons with reduced inhibition and high synaptic plasticity, not on precursor cell proliferation or intermediate progenitor cells *per se*.

Neocortical development is an example of where, in the human brain, a common developmental principle has evolved to greater complexity: a precursor cell population that is only transient in mice and rats became the foundation of the massive expansion and gyration of the neocortex in primates (Fietz et al., 2010). However, the basal progenitor cell that allowed this step at least transiently also exists in mice. With respect to adult neurogenesis, a key difference between rodents and humans might therefore lie in the specific qualitative and quantitative relationship between precursor cell proliferation, a hypothesized non-proliferative waiting state, a period of high synaptic plasticity, and the lasting integration of the new neurons.

The contribution of such highly plastic ‘neurons in waiting’ not only depends on the number of cells but also on the duration of this critical time window of enhanced plasticity (Kempermann, 2012). The period of DCX expression appears to be about a month long in humans as it is in mice, but species might still differ in that respect. In any case, full maturation of newborn neurons might take several months in primates (Kohler et al., 2011), resulting in a heterogeneity of the granule cell population with a relatively large subpopulation of early ‘neurons in waiting’ with delayed final maturation.

Different mammalian species might have developed different solutions to the problem of how to provide a critical population of highly plastic cells to the network. For example, the red fox (*Vulpes vulpes*) has very high numbers of DCX-positive cells but very low levels of proliferation, which is quite different from mice (Amrein and Slomianka, 2010).

The balance between retained neurogenic potential from proliferating progenitor cells or from a reservoir of pre-generated, highly excitable cells might also vary between human individuals (see discussion above and (Spalding et al., 2013)). In addition, this balance is likely to change across the life span. If the duration of the window of plasticity lengthens with age, extremely low numbers of proliferating cells could still contribute to a reservoir of

Author Manuscript
Author Manuscript
Author Manuscript
Author Manuscript

plastic cells that sustain the required functionality. To some extent, this functionality also seems to be additive, in that past neurogenic events also lastingly change the networks (because the new neurons survive for long durations with presumably ‘normal’ levels of synaptic plasticity), so that aged individuals might actually require lower numbers of new neurons.

The process of adult neurogenesis may somewhat parallel what occurs in the female reproductive system of mammals, where all stem cell proliferation that generates the population of egg cells occurs very early in life and further development is delayed. The case of adult neurogenesis might not be as extreme, depending on if the study by Sorrells et al. or Boldrini et al. best reflects the situation, but there is no fundamental need for substantial stem cell proliferation in adult neurogenesis to extend throughout the ever-expanding life span of humans. There might also be a ‘neurogenic menopause,’ in which the potential is used up, and this might indeed contribute to age-related cognitive decline.

Conclusion

Regarding adult hippocampal neurogenesis in humans, many questions remain unanswered. Species differences are interesting and important, and the report by Sorrells et al. reminds us that simple 1:1 translations from animal studies to humans are problematic. But the coincident publication by Boldrini et al., which is more in line with the current body of knowledge briefly summarized in the present article, not only further questions the categorical claim that there is no adult neurogenesis in the human hippocampus but also points out the direction in which this kind of research will develop: towards a more quantitative analysis that aims at relating neurogenesis parameters to other features of plasticity and to behavior in health and disease. There is a clear need for additional ways to study the generation of new neurons in adult humans. A more complete analysis of cell phenotypes and potential differentiation trajectories, by for example single cell RNA-sequencing, is likely to provide valuable information. Methods for following the process of adult neurogenesis *in vivo* would be extremely valuable, particularly for assessing changes under different conditions and in pathology.

Since the serendipitous discovery of adult neurogenesis by Joseph Altman (Altman and Das, 1965) and the heated discussion about ‘Limits of neurogenesis in primates’ (Rakic, 1985) after Fernando Nottebohm’s description of adult neurogenesis in songbirds in the 1980s, the field has come a long way and amassed a more than critical and multifaceted body of evidence supporting the existence of adult neurogenesis in human brains. Human evolution might have found very efficient ways to balance proliferation and the duration of the critical maturation period in order to provide the level of hippocampal plasticity that the individual requires.

Acknowledgments

This research was in part (H.A.C.) supported by the Intramural Research Program of the NIMH.

References

- Abrous DN, Wojtowicz JM. Interaction between Neurogenesis and Hippocampal Memory System: New Vistas. *Cold Spring Harb Perspect Biol*. 2015; 7:a018952. [PubMed: 26032718]
- Altman J, Das GD. Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J Comp Neurol*. 1965; 124:319–335. [PubMed: 5861717]
- Amrein I, Slomianka L. A morphologically distinct granule cell type in the dentate gyrus of the red fox correlates with adult hippocampal neurogenesis. *Brain Research*. 2010; 1328:12–24. [PubMed: 20206610]
- Bao AM, Swaab DF. The art of matching brain tissue from patients and controls for postmortem research. *Handb Clin Neurol*. 2018; 150:197–217. [PubMed: 29496142]
- Bauer S, Patterson PH. The cell cycle-apoptosis connection revisited in the adult brain. *The Journal of Cell Biology*. 2005; 171:641–650. [PubMed: 16291699]
- Boekhoorn K, Joels M, Lucassen PJ. Increased proliferation reflects glial and vascular-associated changes, but not neurogenesis in the presenile Alzheimer hippocampus. *Neurobiol Dis*. 2006; 24:1–14. [PubMed: 16814555]
- Boldrini M, Fulmore CA, Tartt AN, Simeon LR, Pavlova I, Poposka V, Rosoklja GB, Stankov A, Arango V, Dwork AJ, et al. Human hippocampal neurogenesis persists throughout aging. *Cell Stem Cell*. 2018
- Chawana R, Alagaili A, Patzke N, Spoeter MA, Mohammed OB, Kaswera C, Gilissen E, Bennett NC, Ihunwo AO, Manger PR. Microbats appear to have adult hippocampal neurogenesis, but post-capture stress causes a rapid decline in the number of neurons expressing doublecortin. *Neuroscience*. 2014; 277:724–733. [PubMed: 25106130]
- Christian KM, Song H, Ming GL. Functions and dysfunctions of adult hippocampal neurogenesis. *Annu Rev Neurosci*. 2014; 37:243–262. [PubMed: 24905596]
- Curtis MA, Penney EB, Pearson AG, van Roon-Mom WMC, Butterworth NJ, Dragunow M, Connor B, Faull RLM. Increased cell proliferation and neurogenesis in the adult human Huntington's disease brain. *Proceedings of the National Academy of Sciences*. 2003; 100:9023–9027.
- Dennis CV, Suh LS, Rodriguez ML, Kril JJ, Sutherland GT. Human adult neurogenesis across the ages: An immunohistochemical study. *Neuropathol Appl Neurobiol*. 2016; 42:621–638. [PubMed: 27424496]
- Eriksson PS, Perfilieva E, Björk-Eriksson T, Alborn AM, Nordborg C, Peterson DA, Gage FH. Neurogenesis in the adult human hippocampus. *Nat Med*. 1998; 4:1313–1317. [PubMed: 9809557]
- Ernst A, Alkass K, Bernard S, Salehpour M, Perl S, Tisdale J, Possnert G, Druid H, Frisen J. Neurogenesis in the striatum of the adult human brain. *Cell*. 2014; 156:1072–1083. [PubMed: 24561062]
- Fietz SA, Kelava I, Vogt J, Wilsch-Bräuninger M, Stenzel D, Fish JL, Corbeil D, Riehn A, Distler W, Nitsch R, et al. OSVZ progenitors of human and ferret neocortex are epithelial-like and expand by integrin signaling. *Nat Neurosci*. 2010; 13:690–699. [PubMed: 20436478]
- Galán L, Gómez-Pinedo U, Guerrero A, García-Verdugo JM, Matías-Guiu J. Amyotrophic lateral sclerosis modifies progenitor neural proliferation in adult classic neurogenic brain niches. 2017;1–10.
- Ge S, Yang CH, Hsu KS, Ming GL, Song H. A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron*. 2007; 54:559–566. [PubMed: 17521569]
- Germain J, Bruel-Jungerman E, Grannec G, Denis C, Lepousez G, Giros B, Francis F, Nosten-Bertrand M. Doublecortin knockout mice show normal hippocampal-dependent memory despite CA3 lamination defects. *PLoS ONE*. 2013; 8:e74992. [PubMed: 24073232]
- Kempermann G. New neurons for ‘survival of the fittest’. *Nat Rev Neurosci*. 2012; 13:727–736. [PubMed: 22948073]
- Knoth R, Singec I, Ditter M, Pantazis G, Capetian P, Meyer RP, Horvat V, Volk B, Kempermann G. Murine features of neurogenesis in the human Hippocampus across the lifespan from 0 to 100 years. *PLoS ONE*. 2010; 5:e8809. [PubMed: 20126454]

- Kohler SJ, Williams NI, Stanton GB, Cameron JL, Greenough WT. Maturation time of new granule cells in the dentate gyrus of adult macaque monkeys exceeds six months. *Proc Natl Acad Sci USA*. 2011; 108:10326–10331. [PubMed: 21646517]
- Kuhn HG, Eisch AJ, Spalding K, Peterson DA. Detection and Phenotypic Characterization of Adult Neurogenesis. *Cold Spring Harb Perspect Biol*. 2016; 8:a025981. [PubMed: 26931327]
- Liu YWJ, Curtis MA, Gibbons HM, Mee EW, Bergin PS, Teoh HH, Connor B, Dragunow M, Faull RLM. Doublecortin expression in the normal and epileptic adult human brain. *Eur J Neurosci*. 2008; 28:2254–2265. [PubMed: 19046368]
- Lucassen PJ, Oomen CA, Naninck EFG, Fitzsimons CP, van Dam AM, Czéh B, Korosi A. Regulation of Adult Neurogenesis and Plasticity by (Early) Stress, Glucocorticoids, and Inflammation. *Cold Spring Harb Perspect Biol*. 2015; 7:a021303. [PubMed: 26330520]
- Marín-Burgin A, Mongiat LA, Pardi MB, Schinder AF. Unique processing during a period of high excitation/inhibition balance in adult-born neurons. *Science*. 2012; 335:1238–1242. [PubMed: 22282476]
- Mathews KJ, Allen KM, Boerrigter D, Ball H, Shannon Weickert C, Double KL. Evidence for reduced neurogenesis in the aging human hippocampus despite stable stem cell markers. *Aging Cell*. 2017; 16:1195–1199. [PubMed: 28766905]
- Palmer TD, Schwartz PH, Taupin P, Kaspar B, Stein SA, Gage FH. Cell culture. Progenitor cells from human brain after death. *Nature*. 2001; 411:42–43. [PubMed: 11333968]
- Rakic P. Limits of neurogenesis in primates. *Science*. 1985; 227:1054–1056. [PubMed: 3975601]
- Schmidt-Hieber C, Jonas P, Bischofberger J. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature*. 2004; 429:184–187. [PubMed: 15107864]
- Snyder J, Choe J, Clifford M, Jeurling S, Hurley P, Brown A, Kamhi J, Cameron H. Adult-Born Hippocampal Neurons Are More Numerous, Faster Maturing, and More Involved in Behavior in Rats than in Mice. *Journal of Neuroscience*. 2009; 29:14484–14495. [PubMed: 19923282]
- Sorrells SF, Paredes MF, Cebran-Silla A, Sandoval K, Qi D, Kelley KW, James D, Mayer S, Chang J, Auguste KI, et al. Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature*. 2018
- Spalding KL, Bergmann O, Alkass K, Bernard S, Salehpour M, Huttner HB, Boström E, Westerlund I, Vial C, Buchholz BA, et al. Dynamics of hippocampal neurogenesis in adult humans. *Cell*. 2013; 153:1227.




Figure 1. Multiple lines of evidence in support of adult hippocampal neurogenesis in humans
 Data from rodents suggest a particular and specific function for adult-generated neurons of the dentate gyrus, which would be of great relevance to human cognition in health and disease (green box). Three birthdating studies confirm the idea that adult hippocampal neurogenesis exists in humans (dark green box, top), and a much larger set of studies based on ex vivo analyses of precursor cells and marker expression provide supportive evidence (light green box, bottom). Sorrells et al. (2018) have questioned the validity of marker studies (red X), but there is little general support for that claim. The other lines of evidence are untouched by their argumentation.




Figure 2. Consequences of species differences in the course of neurogenesis

Besides methodological considerations, a hypothetical concept of a temporal decoupling of the stages of adult neurogenesis and species differences in marker expression, although largely speculative at this time, might explain part of the discrepancies between rodent and human data. The point is that alternative hypotheses are possible that are consistent with the available data.