

Integrative analysis unveils new functions for the *Drosophila* Cutoff protein in non-coding RNA biogenesis and gene regulation

Yuri Pritykin^{2,3,4}, Tarcisio Brito¹, Trudi Schupbach⁵, Mona Singh^{2,3} and Attilio Pane^{1*}.

¹ Instituto de Ciências Biomédicas (ICB), Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro 21949-902, Brazil. ²The Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544, USA. ³ Department of Computer Science, Princeton University, Princeton, NJ 08544, USA. ⁴ Computational and Systems Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA. ⁵ Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA.

* To whom correspondence should be addressed: Instituto de Ciências Biomédicas (ICB), Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, 21949-902, Brazil. Tel. +55(21)3938-7475, E-mail: apane@icb.ufrj.br. Correspondence may also be addressed to: Department of Molecular Biology, Princeton University, Washington Road, Princeton, NJ 08544, USA. Tel.: þ1 609 258 1365; Fax: þ1 609 258 1547; E-mail: schupbac@princeton.edu.

Keywords

Drosophila; germline; piRNAs; non-coding RNAs; Cutoff

Running title

Shared biogenesis for different classes of non-coding RNAs in *Drosophila*

Pritykin et al

ABSTRACT

Piwi-Interacting RNAs (piRNAs) are central components of the piRNA pathway, which directs transposon silencing and guarantees genome integrity in the germ cells of several metazoans. In *Drosophila*, piRNAs are produced from discrete regions of the genome termed piRNA clusters, whose expression relies on the RDC complex comprised of the core proteins Rhino, Deadlock and Cutoff. To date, the RDC complex has been exclusively implicated in the regulation of the piRNA loci. Here we further elucidate the function of Cutoff and the RDC complex by performing genome-wide CHIP-seq and RNA-seq assays in the *Drosophila* germline and analyzing these data together with other publicly available data sets. In agreement with previous studies, we confirm that Cutoff is involved in the transcriptional regulation of piRNA clusters and in the repression of transposable elements in germ cells. Surprisingly, however, we find that Cutoff is enriched at and affects the expression of other non-coding RNAs, including spliceosomal RNAs (snRNAs) and small nucleolar RNAs (snoRNAs). At least in some instances, Cutoff appears to act at a transcriptional level in concert with Rhino and perhaps Deadlock. Finally, we show that mutations in Cutoff result in the deregulation of hundreds of protein-coding genes in germ cells. Our study uncovers a broader function for the RDC complex in the *Drosophila* germline development.

INTRODUCTION

Non-coding RNAs (ncRNAs) exert essential roles in a variety of developmental events, where they regulate mechanisms ranging from transcription to RNA processing and translation. In recent years, a specific class of small ncRNAs termed Piwi-interacting RNAs or piRNAs have emerged as key components of a defense mechanism that

Pritykin et al

protects the genome against transposable and repetitive sequences. In *Drosophila*, piRNAs are mostly produced from regions of the genome referred to as piRNA clusters (Brennecke et al. 2007). Both in the germline and in the somatic tissues of the ovary, the processing of cluster transcripts leads to the production of mature piRNAs, which interact with the Piwi protein to silence transposable elements (TEs) dispersed in the genome (Brennecke et al. 2007; Malone et al. 2009). In the germ cells, however, two additional members of the Piwi-clade Argonaute family, namely, Aubergine (Aub) and Argonaute3 (Ago3), sustain the production of piRNAs by promoting a feed-forward amplification loop known as the “ping-pong” mechanism (Brennecke et al. 2007; Gunawardane et al. 2007; Li et al. 2009a). piRNA biogenesis in the germline also requires the RNA helicase Vasa and a number of cytoplasmic proteins, which generally accumulate together with Aub and Ago3 in a perinuclear organelle known as the “nuage” (Pek et al. 2012; Xiol et al. 2014; Nishida et al. 2015). While Aub and Ago3 are cytoplasmic germline-specific factors, Piwi is mostly nuclear and is found both in the germ cells and in the somatic follicle cells of the ovary. Piwi bound to antisense piRNAs imposes the transcriptional silencing of TEs by regulating the levels of Histone H3 trimethyl Lysine 9 (H3K9trimeth) and the recruitment of RNA Pol2 at the target element (Sienski et al. 2012).

Genomic regions that have the capacity of generating piRNAs have been identified in distantly related organisms, including mouse and human (Girard et al. 2006; Aravin et al. 2007). In *Drosophila*, piRNA clusters are predominantly composed of fragmented and mostly inactive transposable elements and are commonly divided in two classes. The uni-strand clusters, like the *flamenco* locus (*flam*), are populated by polarized transposon relics and produce piRNAs from one strand of the genome (Dennis et al. 2013; Malone et al. 2009; Mével-Ninio et al. 2007; Sarot et al. 2004). The majority of the

Pritykin et al

piRNA clusters in this species, however, belong to the so-called dual-strand class, where transposon remnants are harbored both on the plus and minus strand of the genome. Paradigmatic of this class is Cluster 1/42AB (Cl1/42AB), which spans roughly 250Kb in the pericentromeric region of the Chromosome II right arm. Like all bidirectional loci, the piRNA pool produced from Cl1/42AB is composed of both sense and antisense piRNAs. Dual-strand cluster expression in germ cells is regulated by the Cutoff (Cuff) protein in a complex with the HP1 variant Rhino (Rhi) (Pane et al. 2011; Klattenhoff et al. 2009). It was recently shown that these two factors also interact with the Deadlock (Del) protein to form the so-called “RDC” complex (Mohn et al. 2014). While Del does not display obvious catalytic domains, Cuff belongs to the Rai1-like family of transcription termination co-factors (Chen et al. 2007). In accordance with this, it was proposed that the RDC complex is essential at piRNA clusters to suppress transcription termination and promote the production of the long piRNA precursor transcripts necessary to generate a full complement of mature piRNAs (Mohn et al. 2014; Chen et al. 2016). Furthermore, the activity of the complex also prevents the splicing of cluster transcripts, which would otherwise eliminate sequences that are potential sources of piRNAs (Zhang et al. 2014; Mohn et al. 2014). This regulatory mechanism requires the activity of the DEAD-box RNA helicase UAP56, an evolutionarily conserved splicing and export factor (Gatfield et al. 2001; Zhang et al. 2012, 2014). UAP56 was shown to co-localize with Rhi in nurse cell nuclear speckles, suppress the splicing of cluster derived transcripts and mediate their transport to the cytoplasmic nuage (Zhang et al. 2012, 2014). Furthermore, UAP56 is a component of the THO/TREX complex, whose component Thoc5 interacts with UAP56 and is necessary for the expression of the piRNA clusters (Hur et al. 2016). While UAP56 has been reported to control the splicing and export of a variety of transcripts, to date the components of the RDC complex have

Pritykin et al

been exclusively connected to the regulation of the piRNA clusters and the biogenesis of piRNAs in *Drosophila* germline tissues.

In the current study, we further investigate the function of the Cuff protein. We perform genome-wide RNA-seq and ChIP-seq assays in *Drosophila* ovaries and combine them with publicly available datasets produced by other laboratories (Mohn et al. 2014; Zhang et al. 2012). In order to obtain a more accurate understanding of the function of Cuff during oogenesis, we used high sequencing depth for each library, adopted 100bp read length protocols, and always produced biological replicates for each assay (Supplementary Table S1). Our integrative computational analysis of these data—consisting of newly generated Cuff ChIP-seq data, RNA Pol2 ChIP-seq and RNA-seq data in wildtype and Cuff mutant ovaries, and analogous existing data for Rhi and Del—confirms that Cuff extensively associates with piRNA cluster sequences and is required for their transcription in germ cells, in agreement with previous reports. Strikingly, our analysis reveals that the Cuff protein is not specific to the piRNA pathway, but also contributes to the expression of several small nuclear RNAs (snRNAs) and small nucleolar RNAs (snoRNAs). snRNAs and snoRNAs have been implicated in various regulatory mechanisms, but they have been best characterized for their role in splicing and ribosomal RNA processing, respectively (Will and Lührmann 2011; Dupuis-Sandoval et al. 2015). We show that Cuff and Rhi ChIP-seq binding profiles often overlap at snRNA and snoRNA loci and that the UAP56 factor associates with snoRNA precursor transcripts. These observations strongly point to a model whereby the RDC complex and the UAP56 factor regulate the biogenesis of different classes of non-coding RNAs in the *Drosophila* germline.

RESULTS

Computational analysis of ChIP-seq data shows significant enrichment of Cuff, Rhi, and Del proteins in non-coding RNA gene loci

Using chromatin immunoprecipitation (ChIP) assays we have previously shown that Cuff is enriched at specific regions within the germline dual-strand piRNA cluster 1/42AB, while it does not bind the uni-strand clusters *flam* and Cluster 2 (Pane et al. 2011). In order to expand our analysis and identify all the Cuff binding sites in the genome, we performed ChIP followed by deep sequencing (ChIP-seq) in ovaries expressing a Cuff-EGFP fusion protein (Pane et al. 2011). DNA fragments from Cuff-EGFP ChIP and control experiments were isolated and used to generate ChIP-seq libraries. We mapped the sequencing reads back to the *Drosophila* genome and pooled all the reads uniquely mapping to the genome. This approach allowed us to identify 937 regions in the *Drosophila* genome that are reproducibly bound by the Cuff protein (IDR threshold 0.05, see Materials and Methods for details). Since Cuff acts in a complex with Rhi and Del, we also analyzed previously published ChIP-seq data for Rhi and Del (Mohn et al. 2014). We assessed the enrichment of Cuff peaks at certain genomic features using a randomization procedure (see Materials and Methods for details). As expected, we could identify a significant enrichment for Cuff at sequences belonging to piRNA clusters (Fig.1A and Supplementary Figure S1-S2). Surprisingly, Cuff appears to strongly associate with other loci that express non-coding RNAs, like snRNA and snoRNA genes. Since Cuff is found in a complex with Rhi, one would expect to observe an enrichment of Rhi at the same genomic features. In accordance with this hypothesis, we can detect a significant overlap between Cuff and Rhi peaks genome-wide and an enrichment of Rhi binding at snRNAs and snoRNAs (Fig.1B and Supplementary Figure S1-S2). The recruitment of the RDC complex at the piRNA clusters relies on the interaction between Rhi and H3K9me3-marked chromatin (Mohn et al. 2014). However,

Pritykin et al

we do not observe a significant enrichment of this histone mark at snRNA and snoRNA loci, suggesting the binding of the RDC complex or some of its components at these ncRNA loci might occur through a different and yet unknown mechanism (Supplementary Figure S2). Consistent with our previous study, we find a robust enrichment of Cuff at Cl1/42AB (Fig.1C), while we did not detect significant binding at the uni-strand *flam* and Cl2. Cuff peaks could not be readily detected at several dual-strand clusters, like the major cluster Cl13, although Rhi peaks are evident (Supplementary Figure S3), and the piRNAs produced from this locus are lost in the *cuff* mutant ovaries (Pane et al. 2011) indicating that Cuff is regulating their expression. This lack of measurable binding might reflect in part the fact that Cuff does not directly interact with the DNA, rather it is recruited by Rhi and Del (Mohn et al. 2014). Intriguingly, our analysis reveals that Cuff and Rhi peaks also overlap at certain snoRNA and snRNA loci. For instance, they are enriched at the snRNA:U4 variant hosted in the intron of the *Atg18b* gene (Fig.1D). These observations pointed to a role for the RDC complex in the regulation of other genomic loci in addition to piRNA clusters. We therefore set to systematically investigate the genome-wide distribution and function of Cuff in germline tissues.

RNA-seq and Pol2 CHIP-seq analysis confirms the role of Cuff in regulating piRNA cluster expression

First, we evaluated the impact of mutations in *cuff* on piRNA cluster expression. We employed RNA-seq assays and compared the expression levels of all the known piRNA loci in wildtype and *cuff* mutant ovaries. For this study, rRNA-depleted total RNA was used to produce cDNA libraries suitable for massive parallel sequencing. We mapped the reads obtained from each assay back to the genome. We then focused on reads that mapped to the genome uniquely. In order to obtain an accurate comparison of different

Pritykin et al

clusters in the wildtype and *cuff* ovaries, we split each piRNA cluster into 500bp fragments, and computed for each of them the fold difference between mutant and wildtype and estimated the differential expression between the two genetic backgrounds (Fig.2A). This approach yielded approximately 9930 unique fragments belonging to the 140 reported piRNA clusters. Despite the depth of our RNA-seq datasets, we could only find uniquely aligned reads for approximately 3730 fragments in at least one of the two genotypes. Of these, 116 displayed significant downregulation (FDR < 0.05) by at least 2-fold in the absence of the Cuff protein, while 86 fragments displayed a significant upregulation (FDR < 0.05) by at least 2-fold in the *cuff* ovaries (false discovery rate calculated using the Benjamini-Hochberg procedure to correct for multiple testing; see Materials and Methods for details). The remaining regions did not show significant changes. We then narrowed our analyses to the major clusters Cl1/42AB and *flam*. We could determine that Cl1/42AB harbors ~130 unique fragments with aligned reads in at least one of the genotypes. None of these fragments are significantly upregulated in the *cuff* mutant, while 30 showed a significant downregulation (at least 2-fold, FDR < 0.05) (Fig. 2B). For the *flam* locus, we could identify ~120 unique fragments with aligned reads in at least one of the genotypes (Fig.2C). Of these, 2 fragments displayed a significant upregulation in the *cuff* mutant (3- to 4-fold), while none of the regions was significantly downregulated. Although limited to a set of unique fragments, our analysis confirms that Cuff is required for the expression of the piRNA cluster Cl1/42AB, while it is not involved in the regulation of the somatic *flam* locus. It is noteworthy that a substantial number of fragments from various clusters displayed increased expression levels in the *cuff* mutant ovaries. It is tempting to speculate that piRNAs might themselves contribute to the expression of the clusters by downregulating certain regions. Alternatively, the transcripts produced

Pritykin et al

from the clusters might, in some cases, fail to be processed into mature piRNAs and might therefore accumulate in germline tissues.

The observation that the steady-state RNA levels of cluster transcripts are sometimes altered in the absence of a functional Cuff protein led us to investigate whether these changes are caused by a transcriptional or post-transcriptional mechanism. To test these alternatives, we attempted to determine the impact of mutations in Cuff on the occupancy of RNA-Pol2 over the piRNA clusters by ChIP-seq assays (Supplementary Figure S4). In accordance with the RNA-seq profiles, Pol2 occupancy does not display uniformly reduced levels at the piRNA clusters in the absence of a functional Cuff protein. Rather, we could identify 49 fragments displaying a substantial increase in Pol2 when we compared the *cuff* mutant versus the wildtype ovaries, while 15 fragments displayed a significant decrease of Pol2 in the *cuff* mutant. Because of the very limited number of regions that displayed significantly altered Pol2 levels in the mutant ovaries, we turned to Pol2 ChIP assays followed by qPCR on selected regions of the dual-strand Cl1/42AB and the uni-strand Cl2 to validate the conclusions of high-throughput data analysis (Fig.2D). As control for this experiment, we analyzed the Pol2 occupancy in wildtype and *cuff* ovaries at the *Pld*, *ry* and *rp49* genes (Fig.2D). This assay revealed a clear loss of Pol2 in the mutant ovaries at regions Cl1A, Cl1A2 and Cl1-32 within the Cl1/42AB cluster. In agreement with our previous results, these were the same regions that we had found to be expressed at a lower level in the absence of a functional Cuff protein (Pane et al. 2011). We also found that region Cl1A6 displayed a clear enrichment of Pol2 in the mutant ovaries, while the levels of Pol2 at region Cl1C appeared unaltered. Similarly, the levels of Pol2 at Cl2A, Cl2C, *Pld*, *ry* and *rp49* were comparable in wildtype and mutant ovaries. These observations strongly suggest that mutations in Cuff perturb the expression pattern of the clusters by affecting the levels of Pol2 along their sequences.

Transposable elements display increased RNA levels and Pol2 occupancy in the *cuff* mutant ovaries

A primary role of the piRNA pathway in *Drosophila* is to silence transposable and repetitive sequences either in germline or in somatic tissues (Senti and Brennecke 2010; Guzzardo et al. 2013). For instance, we reported earlier that *cuff* mutant ovaries display elevated levels of the *Het-A* and *TART* telomere-specific retrotransposons (Chen et al. 2007). We employed our RNA-seq datasets to expand this analysis by comparing the expression levels of all the canonical transposable elements in wildtype versus *cuff* ovaries. *Drosophila* transposons have been divided in 122 families. Our RNA-seq approach allowed us to determine that 40 transposon families are significantly upregulated in the *cuff* mutant (FDR < 0.05), 80 are apparently not significantly affected and only two seem to be significantly downregulated (Fig.2E). Among the transposons that are up-regulated in *cuff*, the values of the fold difference between the mutant and the wildtype vary from approximately 1.4-fold difference for *copia* to more than 120 fold for *gypsy12* and *3S18*. Consistent with our previous study, the *Het-A* transposon appears to be up-regulated by more than 30 fold and *TART-A* by approximately 7 fold in the mutant ovaries. The *TAHRE* element, which is involved in telomere regulation together with *Het-A* and *TART*, is also significantly de-repressed in the absence of Cuff.

We then asked whether the mutations in *cuff* cause transposon deregulation through a transcriptional versus a post-transcriptional mechanism. To answer this question, we compared the RNA-Pol2 occupancy over the transposable elements in wildtype and *cuff* ovaries (Fig.2F). Interestingly, we observed a significant increase in RNA-Pol2 over 19 transposon families in the absence of a functional Cuff protein with fold differences varying from approximately 2-fold to more than 4-fold. These include the *Het-A*, *TART-B*

Pritykin et al

and *TAHRE* elements (Supplementary Figure S5). The *gypsy12* element is the most sensitive to mutations in *cuff* both in terms of transcript levels as well as RNA Pol2 occupancy. When comparing *cuff* over wildtype log fold change values of RNA level with *cuff* over wildtype log fold change values of RNA-Pol2 level across all transposons detectable in both assays, we observed them to be significantly positively correlated (Spearman correlation 0.3615, p-value < 4e-5), which confirms massive transposon deregulation at the transcriptional level.

Intriguingly, Cuff binding events can be detected at several transposon families including the *Het-A*, *TART* and *TAHRE* elements (Supplementary Figure S5). These telomere-specific transposons are at the same time the sources and the targets of piRNAs. Hence, Cuff might control transposon expression not only indirectly, by its role in piRNA cluster regulation and piRNA production, but also directly by associating with sequences within the transposable elements.

Cuff regulates spliceosomal RNA expression

To date, the activity of Cuff and of the RDC complex has been connected only to piRNA cluster regulation in *Drosophila* germ cells. However, our CHIP-seq assays showed that Cuff is enriched at hundreds of positions in the genome including snRNA and snoRNA loci. The *Drosophila* genome harbors 31 snRNAs, which are generally divided into two classes: the U2-type, which catalyzes 99% of the splicing events in eukaryotic cells, and the U12-type (Alioto 2007; Lu and Matera 2014). The major U1, U2, U4, U5 and U6 snRNAs belong to the first class and can display multiple paralogs in the genome, while the U12-type are single copy genes. We find that mutations in Cuff significantly affect the levels of 18 snRNAs belonging to the U2-type, which drop from 2- to more than 8-fold as per RNA-seq in the mutant ovaries, while U12-type snRNAs are not significantly affected (Fig.3A). In order to lend support to our high-throughput analyses, we

Pritykin et al

performed qRT-PCR assays with oligonucleotides specific for the major spliceosomal RNAs in wildtype, *cuff* and *rhi* mutant ovaries (Fig.3B). We could observe a reduction of the U2, U4 and U5 snRNA levels in the *cuff* mutant compared to the wildtype ovaries. In agreement with these observations, a consistent reduction of the snRNA levels can be detected also in the *rhi* mutant ovaries (Fig.3B). The U2 and U5 snRNAs are significantly downregulated by up to 1.8 fold both in the *cuff* and in the *rhi* mutants, thus suggesting that the RDC complex might exert a prominent role in the regulation of these snRNAs.

The comparison of the Pol2 occupancy at these loci by CHIP-seq reveals that in some cases the reduced expression levels are likely caused by loss of Pol2 over the gene region (Fig.3C and Fig.3D). For instance, mutations in *Cuff* cause a partial loss of Pol2 over the genomic region encompassing the snRNAs U4:38Ab, U5:38Aba and U2:38ABb and a concomitant decrease in their RNA levels (Fig.3D). The region spanning these snRNAs is enriched in the *Rhi* protein. Although *Cuff* peaks cannot be clearly detected in this case, the data suggest that sets of neighboring snRNAs can be co-regulated by the RDC complex at a transcriptional level.

Northern blotting on total RNA extracted from ovaries shows a clear accumulation of partial transcripts for the snRNA:U2 in the *cuff* mutant compared to the wildtype (Fig.3E). While we cannot rule out that these fragments might be the result of partial degradation of the snRNA:U2, it is tempting to speculate that they might reflect incomplete transcription of the locus in the absence of *Cuff*. Different from the U2, partial or fragmented transcripts are not detected for U1, U4 and U5 (Fig.3E and data not shown). Nevertheless, our data strongly point to a direct role for *Cuff* and *Rhi* in the regulation of certain snRNA loci in the *Drosophila* germline.

Small nucleolar RNA expression is controlled by the *Cuff* protein

Pritykin et al

SnoRNAs are more numerous and heterogeneous than snRNAs and their functions are still largely unclear. These ncRNAs are generally hosted in the intronic sequences of protein-coding genes and the release of the mature snoRNAs relies on the regulated splicing of the host precursor RNAs. We find that mutations in *cuff* significantly affect 92 of the 281 reported snoRNAs, whose expression levels drop by 2- to 9-fold in the mutant ovaries (Fig. 4A). One remarkable example is represented by an array of snoRNAs residing within the 3'UTR of the Nop60B gene. The expression of the snoRNAs is coupled to the transcription of Nop60B, whereby the usage of alternative transcription termination sites in the 3'UTR of this gene allows the production of transcripts encompassing the snoRNA set (Riccardo et al. 2007). Intriguingly, combining our data with that of Mohn and coworkers (15), we can clearly detect partially overlapping peaks for the Cuff, Rhi and probably Del proteins at the 3' end of Nop60B (Fig.4B). Mutations in *cuff* cause a significant loss of Pol2 over the Nop60B 3'UTR containing the snoRNAs without affecting the Pol2 occupancy along the coding region. This observation is substantiated by the significant downregulation of the snoRNAs belonging to this cluster in the *cuff* ovaries as per RNA-seq, whereas the expression levels of the Nop60B gene are not significantly altered (Fig 4B and 4C). Thus, Cuff, probably in a complex with Rhi and perhaps Del, appears to suppress an alternative transcription termination site at this locus and, in doing so, it couples the expression of the mRNA encoding the Nop60B protein to the production of the snoRNAs in germline tissues.

snoRNAs are mostly found within the intronic sequences of protein-coding genes. In *Drosophila* however, snoRNAs are also generated by specific loci termed Non-coding host genes (Uhg). Uhg genes do not encode for proteins, but serve as carriers of snoRNAs, which are hosted in their intronic sequences (Tycowski and Steitz 2001). The release of mature snoRNAs proceeds through regulated splicing and trimming of the

Pritykin et al

Uhg transcripts. Interestingly, Cuff and Rhi appear to be enriched at Uhg5 (Fig. 4D), thus suggesting that these proteins might directly regulate the expression of the Uhg5 snoRNA set. In order to strengthen these observations, we turned to genome-wide studies independently performed by other laboratories. It was recently shown that the processing and transport of the piRNA cluster transcripts from the nucleus to the cytoplasm is mediated by the DEAD-box helicase UAP56, which is found to colocalize with Rhi in the nurse cell nuclei (Zhang et al. 2012). Zhang and coworkers performed UAP56 RNA immunoprecipitation assays followed by deep sequencing (RIP-seq) and unveiled a significant enrichment of the piRNA precursors in the UAP56 immunoprecipitates (Zhang et al. 2012). We reasoned that if UAP56 is implicated in the regulation of the Uhg loci together with Cuff and Rhi, we would expect to find it associated with the Uhg transcripts. In agreement with this hypothesis, we could determine that Uhg RNAs are strongly associated with the wildtype UAP56 protein, while we did not detect a preferential binding of this factor to protein-coding genes (Fig. 4E). The Uhg5 and Uhg7 transcripts show the strongest enrichment in the wildtype UAP56 protein precipitates (Fig. 4E). Transcripts from the Nop60B also seem to be enriched in UAP56 complexes albeit to a lesser extent, whereas Uhg3 RNAs are not (Fig. 4E). RIP-seq assays in germ cells were performed expressing the tagged wildtype UAP56 or the mutated UAP56^{sz15} variant, where the mutated protein is characterized by an E245K substitution affecting a conserved residue (Zhang et al. 2012). Intriguingly, the binding of UAP56 at Uhg transcripts is significantly compromised by the E245K substitution harbored in UAP56^{sz15} (Fig. 4E). This is particularly evident for Uhg5 and Uhg7 (Fig. 4E) and suggests that this conserved residue is critical for the capacity of UAP56 to specifically recognize and associate with Uhg transcripts. Our study implicates the components of the RDC complex and the UAP56 splicing/export factor in the regulation of snoRNA loci in germline tissues.

RNA-seq and ChIP-seq analysis reveals mutations in *Cuff* affect the expression of hundreds of protein-coding genes

Previous studies reported that mutations in piRNA pathway components do not generally impact the expression levels of protein-coding genes (Klattenhoff et al. 2009; Mohn et al. 2014). We explored our genome-wide datasets to determine whether we reach the same conclusions for the *cuff* mutant. To this aim, we searched our RNA-Pol2 ChIP-seq and RNA-seq datasets for genes that were differentially expressed in wildtype versus *cuff* ovaries. We could identify 526 genes, which displayed a significant upregulation and 313 genes displaying a significant reduction (at least 2-fold, FDR < 0.05) in the absence of *Cuff* (Fig. 5A). The analysis of the Pol2 occupancy at protein-coding genes in wildtype versus *cuff* mutant ovaries suggests that in several cases the observed changes in the expression levels are the result of transcriptional, rather than post-transcriptional events (Fig. 5B). Consistent with this, we observe a significant positive correlation (Spearman correlation 0.25, p-value < 6e-104), when we compare the log fold changes in the steady-state RNA levels and Pol2 occupancy in *cuff* over wt across all genes detectable in both assays. When restricting this comparison to genes with significant over- or under-expression of RNA levels, we observe even stronger correlation (Spearman correlation 0.57, p-value < 2e-50). For instance, the steady-state RNA levels for the uncharacterized CG4415 and CG5111 are approximately 1000-fold and 360-fold higher, respectively, in *cuff* versus wildtype ovaries. Concomitantly, a higher RNA-Pol2 occupancy is observed in the mutant either at the promoter or over the body of the gene, thus pointing to a transcriptional control mechanism (Fig.5C and 5D). The alignment of the RNA-Pol2 ChIP-seq sequence reads to the reference *Drosophila* genome excluded the presence of novel transposon insertions in the vicinity of or within CG4415 and CG5111 in the genome of the *cuff* homozygous mutant flies

Pritykin et al

(Supplementary Bioinformatic Methods). In contrast, the CG43290 lies in close proximity of the *micropia* element on the minus strand of the genome (Fig.5E). This uncharacterized gene is expressed below detectable levels in wildtype ovaries, while it displays elevated expression in the *cuff* mutant. The tandem arrangement suggests the deregulation of the mobile element in *cuff* might be affecting the expression of the downstream CG43290. A similar mechanism is induced upon loss of Piwi in Drosophila OSC cells, where Pol2 spreading from transposon sequences was shown to affect the expression of neighboring loci (Sienski et al. 2012). In addition to genes that are expressed at substantially higher levels, we also find a group of genes with reduced RNA levels in the *cuff* ovaries. For instance, mutations in Cuff appear to cause a loss of RNA-Pol2 at GILT3 gene and a concomitant reduction of the corresponding mRNA levels (Fig.5F). In all the cases we have illustrated so far, Cuff and Rhi peaks are apparently not present within the genes or in the neighboring regions. However, according to our Cuff ChIP-seq assays Cuff is enriched at many protein-coding genes (Fig.1A). In addition, Cuff-bound and Rhi-bound loci display a mild, but significant upregulation in the absence of a functional Cuff protein (Supplementary Fig.S6). Thus we cannot rule out that Cuff and other components of the RDC complex might also contribute directly to the regulation of certain protein-coding loci. The analysis of the CG42854 lends some support to these conclusions (Fig.5G). Rhi binding is apparent at this uncharacterized CG, while Cuff peaks are not detected. Despite the absence of clear Cuff binding however, CG42854 is significantly upregulated in the *cuff* mutant ovaries as per RNA-seq.

In order to confirm the RNA-seq data, we performed RT-PCR assays on total RNA extracted from wildtype and *cuff* mutant ovaries with oligonucleotides specific for CG4415, CG5111, CG43290 and the Rhi-bound CG42854. Oligonucleotides specific for *rp49* were used as a control for this experiment. Consistent with the RNA-seq data, all

Pritykin et al

the genes analyzed are expressed at very low or undetectable levels in wildtype ovaries, while a clear signal is detectable for each of them in the *cuff* mutant ovaries (Fig. 5H). We did not observe significant differences in the signals produced by *rp49* in wildtype and *cuff* ovaries (Fig.5H). The absence of any Cuff, Rhi or Del peaks at or in the vicinity of the CG4415, CG5111 and CG43290 loci suggests that their deregulation is likely an indirect effect of mutations in the Cuff protein. However, the analysis of the CG42854 locus strongly points to a direct role for the RDC complex or at least some of its components in the direct regulation of certain protein-coding genes.

DISCUSSION

In distantly related organisms, piRNAs generated by genomic regions termed piRNA clusters are essential to guarantee transposon silencing and genome integrity during germ cell development. In *Drosophila*, piRNA cluster expression is controlled by the RDC complex, which is centered on the core proteins Cuff, Rhi and Del (Pane et al. 2011; Klattenhoff et al. 2009; Mohn et al. 2014; Le Thomas et al. 2014). To date, this complex has been exclusively implicated in the early steps of piRNA biogenesis in the *Drosophila* germ cells. Surprisingly, our genome-wide analyses unveil a broader function for Cuff in the regulation of different classes of non-coding RNAs. We find that Cuff is not only enriched at piRNA clusters, but it also binds snRNA and snoRNA loci and it is necessary for their expression in germline tissues. Similar to the piRNA clusters, snoRNAs and snRNAs appear to be often enriched for the Rhi protein, suggesting that the underlying regulatory mechanism might be controlled by the RDC complex, rather than by Cuff alone. The role of Cuff in *Drosophila* shares some analogies with the function of the SNAPc complex subunit SNPC-4, which was recently shown to bind and regulate piRNA clusters, snRNAs and snoRNAs in the germline of the nematode worm *C. elegans*. (Kasper et al. 2014).

Pritykin et al

In support of our conclusions, the transcripts produced from snoRNA sources, like the Uhg loci, associate with the UAP56 protein. This splicing/export factor co-localizes with Rhi in nuclear foci and is crucial for the suppression of splicing events at piRNA clusters and the export of the piRNA precursors to the cytoplasmic nuage compartment (Zhang et al. 2012, 2014). Interestingly, the UAP56^{sz} mutant form of the protein is unable to efficiently bind the Uhg transcripts, while the same mutation did not seem to substantially affect the binding to piRNA cluster transcripts (Zhang et al. 2012). It is possible that the amino acid residue that is substituted in the mutated form of UAP56 is essential for the specific interaction between UAP56 and Uhg RNAs. The UAP56 protein was recently shown to interact with Thoc5, a subunit of the THO/TREX complex (Hur et al. 2016). Interestingly, when we analyzed the RNAseq datasets produced by Hur and coworkers (Hur et al. 2016), we found that mutations in Thoc5 result in increased snRNA and snoRNA expression levels (data not shown). The coordinated activity of the RDC complex components, the UAP56 protein and THO/TREX complex might be critical to ensure that different non-coding RNA precursors are expressed and sorted into the appropriate biogenesis pathway.

Recently, Cuff was proposed to prevent transcription termination at piRNA clusters by binding the CAP structure of nascent transcripts and protecting them from degradation (Mohn et al. 2014). Together with the suppression of splicing and polyadenylation events, this mechanism elicits the expression of long piRNA precursors from the clusters, an essential prerequisite for the production of a proper piRNA complement (Mohn et al. 2014; Zhang et al. 2014; Chen et al. 2016). Our study suggests that Cuff, Rhi and Del might also control transcription termination at loci other than the piRNA clusters. This is illustrated by the regulation of the snoRNA set hosted in the 3'UTR of the Nop60B gene, where Cuff, probably in a complex with Rhi and Del, appears to be enriched at the 3'end of the locus and suppress an alternative transcription termination

Pritykin et al

site in the Nop60B gene. This regulatory mechanism promotes the inclusion of the snoRNAs in the nascent Nop60B transcripts and likely couples the expression of the Nop60B protein to the production of the snoRNAs in the germ cells. In addition, our study seems to add new layers of complexity to the repertoire of functions exerted by the Cuff protein. For instance, the loss of Cuff at the snRNA:U4:39B does not appear to affect the overall Pol2 occupancy, although the expression levels of this snRNA are severely diminished in the *cuff* mutant ovaries. Thus, Cuff might act co-transcriptionally to control the production or the stability of the nascent transcript at this locus. Differently, the snRNA variants U4, U5 and U2 hosted in the intron of the *sickle* gene seem to be co-regulated at a transcriptional level. Loss of Cuff causes a clear reduction of the Pol2 levels over the region encompassing these loci. It is conceivable that reduction of the snRNA levels might also contribute to the general downregulation of the snoRNA and piRNA populations given that the biogenesis of both these ncRNA species is based on the regulation or suppression of splicing events.

Importantly, we also find that Cuff appears to control the expression of some protein-coding genes. For example, the uncharacterized gene CG42854 appears to be bound along its length by the Rhi protein and is expressed at very low levels in the wildtype, while its expression is greatly increased in the *cuff* ovaries. This observation reveals that the RDC complex or at least some of its components can function as negative regulators of gene expression in germ cells and that the genomic loci regulated by this complex are not limited to non-coding RNAs, but can include certain protein-coding genes. It is possible that the interaction with additional factors together with inherent structural motifs in the nascent RNAs or *cis*-acting elements at the regulated loci might exert an essential role in defining the specific activity of Cuff. This functional flexibility is probably crucial at piRNA clusters to bypass *cis*-regulatory elements (i.e. transcription termination and splice sites) carried by the transposon sequences and

Pritykin et al

guarantee the production of long piRNA precursors that can be efficiently converted into a full set of piRNAs.

To date, mutations in piRNA pathway components have not been found to affect the transcript levels of protein-coding genes in the *Drosophila* germ cells. However, we could identify a few hundred loci with dramatically altered steady state RNA levels in the *cuff* mutant ovaries. In some cases, the Pol2 ChIP-seq assays support a model whereby loss of Cuff causes a change in the transcriptional activity at certain loci. For example, higher expression levels of CG4415 and CG5111 are consistent with a significant accumulation of Pol2 at their promoter regions and over the gene body in the *cuff* mutant. The absence of Cuff peaks at these CGs as well as at other genes suggests that Cuff is only indirectly involved in their transcriptional regulation. We and others have reported that the loss of the piRNA pathway activities leads to the activation of a meiotic checkpoint system, centered on the Checkpoint kinase Chk2, in germline tissues (Chen et al. 2007; Klattenhoff et al. 2007). This checkpoint senses the DNA damage induced by massive transposon mobilization and translates it into the loss of germ and stem cells, the defective assembly of the oocyte nucleus (i.e. karyosome phenotype) and the disruption of the Dorsal-Ventral axis of the egg and future embryo, which characterize *cuff* as well as other piRNA pathway mutants. Although the details of the signaling mechanism are not known, it is conceivable that Chk2 might act by regulating the expression of yet-unknown downstream genes. In this scenario, the deregulation of certain protein-coding loci observed in *cuff* mutant ovaries might at least in part depend upon the activation of checkpoint mechanisms in the germline. In accordance with this, we find that the expression levels of CG4415 and CG5111, which are dramatically increased in the *cuff* mutant, are restored to wildtype levels in ovaries doubly mutant for *cuff* and *chk2* (A.P. unpublished observation). It is therefore tempting to speculate

Pritykin et al

that these CGs might belong to a set of genes that form a Chk2-mediated cellular response to the transposon-induced DNA damage.

Our data strongly implicate Cuff and the RDC complex in the biogenesis of piRNAs, snRNAs and snoRNAs and reveal a previously unknown degree of crosstalk between different classes of ncRNAs in the *Drosophila* germline. In line with our findings, Klenov and coworkers have recently shown that loss of nuclear Piwi affects the levels of certain spliceosomal RNAs, including the snRNA:U2:Aba, snRNA:U5:38ABa and snRNA:U4:39B, that we find to be downregulated in *cuff* ovaries (Klenov et al. 2014). The discovery that the initial steps in the production of different classes of non-coding RNAs relies on a shared set of proteins will help shed light on the biogenesis of piRNAs, snRNAs and snoRNAs in the *Drosophila* germ cells.

MATERIALS AND METHODS

Fly stocks

The *cuff*^{WM25} allele used in this study was isolated in a screening for female sterile mutations (Schüpbach and Wieschaus 1989). The *rhi*^{KG00910} allele originates from a P-element insertion in the *rhi* locus (Volpe et al. 2001) (Bloomington Stock Center). *rhi*^{KG00910} was used in combination with the deficiency Df(2R)Exel17149 (Bloomington Stock Center), that spans the *rhi* gene. Heterozygous *cuff*^{WM25}/CyO or *rhi*^{KG00910}/CyO flies were used as wildtype control.

RNA-seq and ChIP-seq assays

RNA-seq and ChIP-seq assays were performed in triplicate and for each biological replicate we dissected ovaries from approximately 30 female flies (2-4 days old). Total RNA was extracted using Trizol reagents (Life Technologies) and treated with turbo

Pritykin et al

DNase-free kit (Ambion) to eliminate traces of genomic DNA. DNA-free total RNA was subjected to two rounds of rRNA depletion using the Ribo-Zero rRNA removal kit (Epicentre) as per the manufacturer's instructions. The RNA-seq library preparation was obtained with the ScriptSeq v2 kit (Epicentre) and the libraries were sequenced on the Illumina platforms.

The conditions and the oligonucleotides used for the Pol2 Chromatin Immunoprecipitation followed by qPCR (ChIP-qPCR) have been previously described (Pane et al. 2011; Blythe et al. 2009). Pol2 ChIP-qPCR assays were performed using anti Pol2 antibodies (clone 8WG16, Millipore). Pol2 enrichment at specific regions of the piRNA cluster 1/42AB, Cl2, *ry*, Pld and *rp49* was calculated as % of the Input chromatin and displayed as fold change [$\log_2(\text{cuff}/\text{wt})$]. The same ChIP protocol was employed for the ChIP-seq assays, where chromatin fragments were used to generate DNA libraries with the ChIP-seq DNA Sample Prep kit (Illumina). The Cuff-EGFP expressing transgenic line, GFP antibodies and the conditions used for the Cuff ChIP-seq experiment were previously described (Pane et al. 2011; Blythe et al. 2009).

Bioinformatic analyses

For ChIP-seq analysis, reads were aligned uniquely to the genome using Bowtie (Langmead and Salzberg 2012). Cuff and Rhi peaks were called using MACS2 (Zhang et al. 2008), and reproducible peaks were detected using IDR (Zhang et al. 2008) threshold 0.05 (for Rhi, publicly available ChIP-seq data from Mohn et al. 2014 was used). Peaks were randomly shuffled in the genome (preserving chromosome and peak length) in order to assess significance of overlap of peaks with genomic features. For visualization in USCS Genome Browser (Kent et al. 2002), bigWig files with genome-wide background subtracted ChIP signal were produced using bedtools (Quinlan and Hall 2010). For RNA-seq analysis, reads were aligned uniquely to the genome using Tophat (Kim et al.

Pritykin et al

2013), and reads aligned to each genomic feature of interest (protein coding gene, snRNA, snoRNA, piRNA cluster 500bp fragment) counted with custom scripts utilizing python interface to samtools (package pysam) (Li et al. 2009b). Differential expression analysis was run using DESeq (Anders and Huber 2010). The Benjamini-Hochberg procedure was used to calculate false discovery rates (FDR) for multiple hypotheses testing correction. Genomic features with zero read count in all replicates were excluded from the analysis. Differential Pol2 occupancy was analyzed in the same manner. Analysis for canonical transposons was run similarly with the exception that reads aligned multiple times to the same transposon were also taken into account. We also used publicly available RIP-seq data (Zhang et al. Cell 2012) to analyze RNA binding profile for UAP56. For this we employed bedGraph files with genome-wide normalized RIP-seq signal at single-nucleotide resolution produced by the authors and available at Gene Expression Omnibus. Detailed description of the bioinformatic analyses is provided in the Supplementary Data. Total number of raw and aligned reads for each library is shown in Supplementary Table 1.

Northern blotting, RT-PCR and qRT-PCR

Fifteen micrograms total RNA extracted from *cuff^{WM25}* and *cuff^{WM25}/Cyo* flies were separated on 6% Urea gels (Life Technologies) using SureLock System (Life Technologies). The RNA was then transferred onto Nylon membranes (Genescreen Plus) and hybridized with P³²-labeled antisense oligonucleotide probes corresponding to sequences in snRNA:U1 and snRNA:U2. Hybridization conditions were previously described (Pane et al. 2007). Probe sequences are reported in the Supplementary Table 2.

Rt-PCR and qRt-PCR reagents and condition and oligonucleotides specific for *rp49*, *Pld*, *ry* and for selected regions within piRNA clusters have been previously described (Pane

Pritykin et al

et al. 2011; Chen et al. 2007). Oligonucleotides specific for protein-coding genes and for snRNAs, which were designed for the current study, are described in Supplementary Table 2.

ACKNOWLEDGEMENTS

We would like to thank Lance Parsons, Donna Storton and Jessica Buckles in the Microarray Facility at Princeton University for help with RNA-seq and ChIP-seq library sequencing; Gail Barcelo, Natalia Ferreira and Aline de Graça Sampaio for assistance with molecular techniques. We are grateful to Paul Schedl, Michael Sammeth and Helena Araujo for critical reading of the manuscript and members of the Schupbach, Singh, Wieschaus and Araujo laboratories for helpful discussions and suggestions.

FUNDING

This work was supported in part by US Public Health Service Grant NIH-R01 GM077620 (TS and AP) and NIH-R01 GM076275 (MS and YP), by NSF 1458457 (MS), by the Howard Hughes Medical Institute (TS), by the CNPq Ciencia Sem Fronteiras Program 410594/2012-2 (AP), by the INCT (AP) and by FAPERJ 218071 (AP).

REFERENCES

- Alioto TS. 2007. U12DB: A database of orthologous U12-type spliceosomal introns. *Nucleic Acids Res* **35**.
- Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol* **11**: R106.
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3218662&tool=pmcentrez&rendertype=abstract>.
- Aravin AA, Sachidanandam R, Girard A, Fejes-Toth K, Hannon GJ. 2007. Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science* **316**: 744–747.

Pritykin *et al*

- Blythe SA, Reid CD, Kessler DS, Klein PS. 2009. Chromatin immunoprecipitation in early *Xenopus laevis* embryos. *Dev Dyn* **238**: 1422–1432.
- Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ. 2007. Discrete Small RNA-Generating Loci as Master Regulators of Transposon Activity in *Drosophila*. *Cell* **128**: 1089–1103.
- Chen Y, Pane A, Schüpbach T. 2007. cutoff and aubergine Mutations Result in Retrotransposon Upregulation and Checkpoint Activation in *Drosophila*. *Curr Biol* **17**: 637–642.
- Chen YCA, Stuwe E, Luo Y, Ninova M, Le Thomas A, Rozhavskaya E, Li S, Vempati S, Laver JD, Patel DJ, et al. 2016. Cutoff Suppresses RNA Polymerase II Termination to Ensure Expression of piRNA Precursors. *Mol Cell* **63**: 97–109.
- Dennis C, Zanni V, Brasslet E, Eymery A, Zhang L, Mteirek R, Jensen S, Rong YS, Vaury C. 2013. “Dot COM”, a Nuclear Transit Center for the Primary piRNA Pathway in *Drosophila*. *PLoS One* **8**.
- Dupuis-Sandoval F, Poirier M, Scott MS. 2015. The emerging landscape of small nucleolar RNAs in cell biology. *Wiley Interdiscip Rev RNA* **6**: 381–397.
- Gatfield D, Le Hir H, Schmitt C, Braun IC, Köcher T, Wilm M, Izaurralde E. 2001. The DEXH/D box protein HEL/UAP56 is essential for mRNA nuclear export in *Drosophila*. *Curr Biol* **11**: 1716–1721.
- Girard A, Sachidanandam R, Hannon GJ, Carmell MA. 2006. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature* **442**: 199–202.
- Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, Nagami T, Siomi H, Siomi MC. 2007. A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science* **315**: 1587–1590.
- Guzzardo PM, Muerdter F, Hannon GJ. 2013. The piRNA pathway in flies: Highlights and future directions. *Curr Opin Genet Dev* **23**: 44–52.
- Hur JK, Luo Y, Moon S, Ninova M, Marinov GK, Chung YD, Aravin AA. 2016. Splicing-independent loading of TREX on nascent RNA is required for efficient expression of dual-strand piRNA clusters in *Drosophila*. *Genes Dev* **30**: 840–855.
- Kasper DM, Wang G, Gardner KE, Johnstone TG, Reinke V. 2014. The *C. elegans* SNAPc component SNPC-4 coats piRNA domains and is globally required for piRNA abundance. *Dev Cell* **31**: 145–158.
- Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler a. D. 2002. The Human Genome Browser at UCSC. *Genome Res* **12**: 996–1006.

Pritykin et al

<http://www.ncbi.nlm.nih.gov/pubmed/186604>\n<http://www.genome.org/cgi/doi/10.1101/gr.229102>.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol* **14**: R36.

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4053844&tool=pmcentrez&rendertype=abstract>.

Klattenhoff C, Bratu DP, McGinnis-Schultz N, Koppetsch BS, Cook HA, Theurkauf WE. 2007. Drosophila rasiRNA Pathway Mutations Disrupt Embryonic Axis Specification through Activation of an ATR/Chk2 DNA Damage Response. *Dev Cell* **12**: 45–55.

Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, Zhang F, Schultz N, Koppetsch BS, Nowosielska A, et al. 2009. The Drosophila HP1 Homolog Rhino Is Required for Transposon Silencing and piRNA Production by Dual-Strand Clusters. *Cell* **138**: 1137–1149.

Klenov MS, Lavrov SA, Korbut AP, Stolyarenko AD, Yakushev EY, Reuter M, Pillai RS, Gvozdev VA. 2014. Impact of nuclear Piwi elimination on chromatin state in Drosophila melanogaster ovaries. *Nucleic Acids Res* **42**: 6208–6218.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357–359.

Le Thomas A, Stuwe E, Li S, Du J, Marinov G, Rozhkov N, Chen YCA, Luo Y, Sachidanandam R, Toth KF, et al. 2014. Transgenerationally inherited piRNAs trigger piRNA biogenesis by changing the chromatin of piRNA clusters and inducing precursor processing. *Genes Dev* **28**: 1667–1680.

Li C, Vagin V V., Lee S, Xu J, Ma S, Xi H, Seitz H, Horwich MD, Syrzycka M, Honda BM, et al. 2009a. Collapse of Germline piRNAs in the Absence of Argonaute3 Reveals Somatic piRNAs in Flies. *Cell* **137**: 509–521.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009b. The Sequence Alignment / Map (SAM) Format and SAMtools 1000 Genome Project Data Processing Subgroup. *Bioinformatics* **25**: 1–2.
<http://www.ncbi.nlm.nih.gov/pubmed/19505943>.

Lu Z, Matera AG. 2014. Developmental analysis of spliceosomal snRNA isoform expression. *G3 (Bethesda)* **5**: 103–10.
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4291461&tool=pmcentrez&rendertype=abstract>.

Pritykin et al

- Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R, Hannon GJ. 2009. Specialized piRNA Pathways Act in Germline and Somatic Tissues of the *Drosophila* Ovary. *Cell* **137**: 522–535.
- Mével-Ninio M, Pelisson A, Kinder J, Campos AR, Bucheton A. 2007. The flamenco locus controls the gypsy and ZAM retroviruses and is required for *Drosophila* oogenesis. *Genetics* **175**: 1615–1624.
- Mohn F, Sienski G, Handler D, Brennecke J. 2014. The Rhino-Deadlock-Cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in *Drosophila*. *Cell* **157**: 1364–1379.
- Nishida KM, Iwasaki YW, Murota Y, Nagao A, Mannen T, Kato Y, Siomi H, Siomi MC. 2015. Respective functions of two distinct siwi complexes assembled during PIWI-interacting RNA biogenesis in bombyx germ cells. *Cell Rep* **10**: 193–203.
- Pane A, Jiang P, Zhao DY, Singh M, Schübach T. 2011. The Cutoff protein regulates piRNA cluster expression and piRNA production in the *Drosophila* germline. *EMBO J* **30**: 4601–4615.
- Pane A, Wehr K, Schübach T. 2007. zucchini and squash Encode Two Putative Nucleases Required for rasiRNA Production in the *Drosophila* Germline. *Dev Cell* **12**: 851–862.
- Pek JW, Patil VS, Kai T. 2012. piRNA pathway and the potential processing site, the nuage, in the *Drosophila* germline. *Dev Growth Differ* **54**: 66–77.
- Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–842.
- Riccardo S, Tortoriello G, Giordano E, Turano M, Furia M. 2007. The coding/non-coding overlapping architecture of the gene encoding the *Drosophila* pseudouridine synthase. *BMC Mol Biol* **8**: 15.
- Sarot E, Payen-Groschêne G, Bucheton A, Péliesson A. 2004. Evidence for a piwi-Dependent RNA Silencing of the gypsy Endogenous Retrovirus by the *Drosophila melanogaster* flamenco Gene. *Genetics* **166**: 1313–1321.
- Schübach T, Wieschaus E. 1989. Female sterile mutations on the second chromosome of *Drosophila melanogaster*. I. Maternal effect mutations. *Genetics* **121**: 101–117.
- Senti KA, Brennecke J. 2010. The piRNA pathway: A fly's perspective on the guardian of the genome. *Trends Genet* **26**: 499–509.
- Sienski G, Dönertas D, Brennecke J. 2012. Transcriptional silencing of transposons by Piwi and maelstrom and its impact on chromatin state and gene expression. *Cell*

Pritykin et al

151: 964–980.

Tycowski KT, Steitz JA. 2001. Non-coding snoRNA host genes in *Drosophila*: expression strategies for modification guide snoRNAs. *Eur J Cell Biol* **80**: 119–125.

Volpe AM, Horowitz H, Grafer CM, Jackson SM, Berg CA. 2001. *Drosophila* rhino encodes a female-specific chromo-domain protein that affects chromosome structure and egg polarity. *Genetics* **159**: 1117–1134.

Will CL, Lührmann R. 2011. Spliceosome structure and function. *Cold Spring Harb Perspect Biol* **3**: 1–2.

Xiol J, Spinelli P, Laussmann MA, Homolka D, Yang Z, Cora E, Couté Y, Conn S, Kadlec J, Sachidanandam R, et al. 2014. RNA clamping by Vasa assembles a piRNA amplifier complex on transposon transcripts. *Cell* **157**: 1698–1711.

Zhang F, Wang J, Xu J, Zhang Z, Koppetsch BS, Schultz N, Vreven T, Meignin C, Davis I, Zamore PD, et al. 2012. UAP56 couples piRNA clusters to the perinuclear transposon silencing machinery. *Cell* **151**: 871–884.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers RM, Brown M, Li W, et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**: R137.

Zhang Z, Wang J, Schultz N, Zhang F, Parhad SS, Tu S, Vreven T, Zamore PD, Weng Z, Theurkauf WE. 2014. The HP1 homolog Rhino anchors a nuclear complex that suppresses piRNA precursor splicing. *Cell* **157**: 1353–1363.

FIGURE LEGENDS

Figure 1. Genome-wide analysis of Cuff activity in *Drosophila* germ cells. (A-B) Association of Cuff and Rhi binding sites with prominent genomic features. A dot (Cuff, red; Rhi, black) shows the number of peaks overlapping with a particular genomic feature (e.g. exons or snRNAs). A boxplot shows the distribution, from 1000 random trials, of the number of overlaps when peaks are randomly shuffled while preserving their size and chromosome (see Supplementary Data for details). Single asterisk indicates $p < 0.05$, double asterisks indicate $p < 0.001$. (C-D) Del (blue), Rhi (black) and Cuff (red) background subtracted ChIP-seq profiles along the Cl1/42AB cluster and the

Pritykin et al

genomic region encompassing a snRNA:U4 gene. Loci reproducibly bound by Cuff (red) and Rhi (black) across replicates are marked with solid lines in separate tracks (not reported for Del due to lack of biological replicates). The black box highlights the Cuff and Rhi peaks at the snRNA:U4 locus. The Rhi and Del ChIP-seq datasets were produced by Mohn and coworkers (Mohn et al. 2014).

Figure 2. Mutations in Cuff alter the transcription of piRNA clusters and transposable elements. (A) Steady state RNA levels at piRNA clusters. Scatter plot displays 500bp fragments of piRNA clusters that are upregulated (blue dots), downregulated (red dots) or not significantly affected (yellow dots) in the *cuff* ovaries (normalized read counts). (B-C) Barplot displaying the fold difference between *cuff* and wildtype ovaries [\log_2 (*cuff/wt*)] at 500bp fragments of Cl1/42AB and *flam*. Fragments with significantly higher (blue bars) and lower (red bars) expression levels in the *cuff* mutant are shown. Yellow bars indicate regions that were not significantly affected by mutations in Cuff. (D) RNA Pol2 occupancy was analyzed by ChIP-qPCR at specific positions at Cl1/42AB, Cl2 and the control *Pld*, *ry* and *rp49* genes. Values are reported as fold difference between *cuff* and wt [\log_2 (*cuff/wt*)]. Error bars represent standard deviation. (E) Scatter plot displaying the distribution of the RNA-seq reads mapping to canonical transposon families in *cuff* versus wildtype ovaries. Transposons that are upregulated (blue dots), downregulated (red dots) or nor significantly altered (yellow dots) in the *cuff* ovaries are reported. Shown on the x- and y-axes are consensus normalized read coverage values computed over replicates. (F) Scatter plot displaying the distribution of RNA Pol2 ChIP-seq reads mapping to canonical transposons in *cuff* versus wildtype ovaries. Transposons showing increased (blue dots), reduced (red dots) or unaltered (yellow dots) Pol2 occupancy in the *cuff* mutant are indicated. Shown on the x- and y-

Pritykin et al

axes of scatter plots are consensus normalized read coverage values computed over replicates.

Figure 3. Cuff controls the expression of spliceosomal RNAs. (A) Barplot of RNA-seq fold difference between wildtype and *cuff* ovaries for the small nuclear RNAs [\log_2 (*cuff*/wt)]. snRNAs with significantly lower (red bars) expression levels in the *cuff* mutant are shown. Yellow bars indicate snRNAs that are not significantly affected by mutations in Cuff. (B) Impact of mutations in *cuff* and *rhi* on the expression levels of the snRNAs U1, U2, U4 and U5 as per qRt-PCR. Y-axis displays the fold difference [\log_2 (FC)] between *cuff* and wt (blue bars) and between *rhi* and wt (red bars). Error bars represent standard deviation. (C) RNA Pol2 ChIP-seq levels at small nuclear RNAs. Barplot shows snRNAs characterized by reduced (red) or not significantly changed (yellow) RNA Pol2 occupancy in *cuff* versus wildtype (normalized read counts) [\log_2 (*cuff*/wt)]. (D) Del (blue), Rhi (black), Cuff (red), RNA Pol2 in wildtype (green), RNA Pol2 in *cuff* mutant (magenta) ChIP-seq profiles along selected snRNAs. Loci reproducibly bound by Rhi are marked with solid line in separate track (not reported for Del due to lack of biological replicates). RNA-seq profiles in wildtype (dark green) and *cuff* mutant (purple) are also shown. Rhi and Del ChIP-seq datasets were produced by Mohn and coworkers (15). (E) Northern blotting analysis of total RNA extracted from wildtype and *cuff* mutant ovaries. Oligonucleotide probes specific to sequences in the snRNA:U1 and snRNA:U2 were used to analyze snRNA expression levels in wildtype and *cuff* ovaries. Red asterisks mark partial transcript of the snRNA:U2, which accumulate in the *cuff* mutant background.

Figure 4. Small nucleolar RNAs are downregulated in *cuff* mutant ovaries. (A) Scatter plot displaying the snoRNAs that are upregulated (blue), downregulated (red) and not

Pritykin et al

significantly affected (yellow) in the *cuff* mutant as per RNA-seq. Expression levels of protein-coding genes are displayed in the background (black). Shown on x- and y-axes are consensus normalized read coverage values computed over replicates. (B) Del (blue), Rhi (black) and Cuff (red), RNA Pol2 in wildtype (green) and RNA Pol2 in *cuff* mutant (magenta) ChIP-seq profiles along Nop60B locus. Loci reproducibly bound by Cuff (red) and Rhi (black) are marked with solid lines in separate tracks (not reported for Del due to lack of biological replicates). RNA-seq profiles in wildtype (dark green) and *cuff* mutant (purple) are also shown. Black frame box highlights the snoRNA cluster in the Nop60B 3'UTR and downstream region that is bound by the RDC complex components. (C) Barplot of the fold difference between wildtype and *cuff* ovaries for the Nop60B coding region and for the snoRNAs hosted within its 3'UTR as per RNA-seq [\log_2 (*cuff*/wt)]. (D) Del (blue), Rhi (black) and Cuff (red) ChIP-seq profiles along the Uhg5 gene. Regions reproducibly bound by Cuff (red) and Rhi (black) as per ChIP-seq are marked with solid lines. (E) UAP56 RIP-seq at Uhg transcripts. Barplot shows the fold enrichment of the RIP-seq signal over corresponding input for the exons of the Uhg genes. Barplot for the wildtype UAP56 (UAP), the mutated UAP56^{sz15} variant (sz) and for the control (w1) are shown. Only the last seven exons were used for Uhg6 (Nop60B), and only exons not overlapping Rpl23A locus were used for Uhg7, to avoid contamination from protein-coding sequence. Median of the values for all genes is shown in the last column. This analysis was performed based on datasets from Zhang and coworkers (Zhang et al. 2012).

Figure 5. Protein-coding genes are differentially expressed in the absence of a functional Cuff protein. (A) Scatter plot displaying the impact of mutations in *cuff* on the expression levels of protein-coding genes as per RNA-seq. Genes that are upregulated (blue dots), downregulated (red dots) and not significantly affected (yellow dots) in the

Pritykin et al

cuff mutant are displayed. Shown on x- and y-axes are consensus normalized read counts over replicates. (B) Scatter plot displaying changes in RNA Pol2 occupancy at protein-coding genes as per ChIP-seq assays. Genes characterized by increased (blue dots) and decreased (red dots) RNA Pol2 occupancy in the *cuff* mutant are reported. Yellow dots indicate genes that display apparently unaffected RNA Pol2 occupancy. Shown on x- and y-axes are normalized consensus read counts over replicates. (C-F) RNA Pol2 in wildtype (green) and RNA Pol2 in *cuff* mutant (magenta) ChIP-seq profiles along selected protein-coding genes. RNA-seq profiles in wildtype (dark green) and *cuff* mutant (purple) are also shown. (G) Rhi (black), RNA Pol2 in wildtype (green) and RNA Pol2 in *cuff* mutant (magenta) ChIP-seq profiles along CG42854. Reproducible Rhi binding is shown by a solid line in a separate track. RNA-seq profiles in wildtype (dark green) and *cuff* mutant (purple) are also shown. (H) RT-PCR in wildtype and *cuff* mutant ovaries with primers specific for the CG4415, CG5111, CG43290, CG42854 and the control *rp49*.

Fig1




Fig2




Fig3




Fig4





Fig5





RNA

A PUBLICATION OF THE RNA SOCIETY

Integrative analysis unveils new functions for the *Drosophila* Cutoff protein in non-coding RNA biogenesis and gene regulation

Yuri Pritykin, Tarcisio Brito, Trudi Schupbach, et al.

RNA published online April 18, 2017

Supplemental Material <http://rnajournal.cshlp.org/content/suppl/2017/04/18/rna.058594.116.DC1>

P<P Published online April 18, 2017 in advance of the print journal.

Accepted Manuscript Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.

Creative Commons License This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>
