



CrossMark
click for updates

Cite this: *Chem. Sci.*, 2014, 5, 4173

Received 19th July 2014
Accepted 4th August 2014

DOI: 10.1039/c4sc02155j

www.rsc.org/chemicalscience

Amine α -heteroarylation *via* photoredox catalysis: a homolytic aromatic substitution pathway†

Christopher K. Prier and David W. C. MacMillan*

The direct α -heteroarylation of tertiary amines has been accomplished *via* photoredox catalysis to generate valuable benzylic amine pharmacophores. A variety of five- and six-membered chloroheteroarenes are shown to function as viable coupling partners for the α -arylation of a diverse range of cyclic and acyclic amines. Evidence is provided for a homolytic aromatic substitution mechanism, in which a catalytically-generated α -amino radical undergoes direct addition to an electrophilic chloroarene.


Introduction

Heterocycles and heteroaromatics are ubiquitous components of biological systems and pharmaceutical research. These invaluable structural motifs are broadly employed in the development of medicinal agents due to their capacity to (i) drive therapeutic potency, (ii) lower drug lipophilicity, (iii) increase aqueous solubility, and (iv) reduce the inhibition of cytochrome P450s.¹ Indeed, at the present time heteroaromatics feature in approximately half of the top 200 pharmaceuticals sold worldwide.² It is not surprising therefore that coupling reactions which generically connect heterocyclic moieties to other molecular fragments are now mainstay transformations in medicinal agent synthesis. For example, couplings between heterocycles and amines *via* C–N bond-forming reactions, either *via* nucleophilic aromatic substitution (S_NAr)³ or transition metal-catalyzed approaches,⁴ have been extensively studied and applied (Fig. 1, eqn (1)).⁵ In contrast, fragment coupling methods that unite heteroaromatic units with the amine α -carbon position (Fig. 1, eqn (2)) are relatively unknown, despite the opportunity to access medicinal agents with similar physiological properties and with proven clinical utility (*e.g.* Veliparib,⁶ Pradaxa,⁷ and Benthiaivalcarb⁸). While important strategies for amine α -arylation have been reported by Dieter,⁹ Campos,¹⁰ Nakamura,¹¹ and others,¹² a generic yet mild α -arylation protocol remains largely elusive.

Visible light photoredox catalysis has emerged in recent years as a powerful platform for the development of new reactions in organic synthesis.¹³ In this context, we have recently demonstrated that photoredox catalysis can be employed to mediate the α -arylation of amines using electron-deficient benzonitrile coupling partners.¹⁴ In this protocol, activation of


the amine α -C–H bond is achieved *via* single-electron oxidation followed by α -deprotonation to generate the critical α -amino radical **1** (Scheme 1).¹⁵ Concomitant single-electron reduction of an accompanying benzonitrile provides a stabilized radical anion, which undergoes selective hetero radical–radical coupling with the neutral radical species **1** to forge the desired α -amino arylation products. Taking inspiration from the Minisci reaction,¹⁶ we recently questioned whether a photoredox-based amine α -arylation could be achieved by the addition of an α -amino radical **1** to a ground-state, neutral arene *via* a homolytic aromatic substitution pathway.^{17–19} This approach would constitute an alternative catalytic mechanism that would

Amines and Heteroarenes: Ubiquitous Components of Drugs




■ Heterocycles: drug-like physical properties, present in ~50% of top 200 drugs

C–N bond-forming amine–heterocycle coupling: wide utility




C–C bond-forming amine–heterocycle coupling: this work



Merck Center for Catalysis at Princeton University, Princeton, New Jersey, 08544, USA.
E-mail: dmacmill@princeton.edu; Fax: +1 609 2585922; Tel: +1 609 2582254

† Electronic supplementary information (ESI) available: Experimental procedures, structural proofs, and spectral data for all new compounds are provided (97 pages) (PDF). See DOI: 10.1039/c4sc02155j

Fig. 1 Significance and approaches to amine–heteroaromatic coupling.



Scheme 1 Proposed arylation via homolytic aromatic substitution.

obviate the need for cyanoarenes and the accompanying radical anion pathway. Herein, we report the successful execution of this plan and describe a new coupling mechanism that employs chloroheteroarenes for the direct α -arylation of amines.

Design plan

As shown in Scheme 1, our proposed mechanism called for a sufficiently oxidizing photocatalyst, *e.g.* iridium complex $\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ ($\text{ppy} = 2$ -phenylpyridine, $\text{dtbbpy} = 4,4'$ -di-*tert*-butyl-2,2'-bipyridine, $E_{1/2}^{\text{Ir(III)/Ir(II)}} = +0.66 \text{ V vs. SCE}$)²⁰ to oxidize the amine substrate *via* a single electron transfer (SET) mechanism ($E_{1/2}^{\text{red}} = +0.71 \text{ V vs. SCE}$ for *N,N*-dimethylaniline).²¹ This

electron transfer step would lead to a reduced iridium complex $\text{Ir}^{\text{II}}(\text{ppy})_2(\text{dtbbpy})$ with concomitant formation of the amine radical cation 2, a species that may be deprotonated at the α -C-H position to generate the α -amino radical 1.²² Addition of this neutral radical species 1 to a highly electrophilic heteroarene 3 *via* homolytic aromatic substitution was expected to provide the radical σ -complex 4, which may readily undergo single-electron reduction ($E_{1/2}^{\text{red}} = -0.26 \text{ V vs. SCE}$ for the hydroxycyclohexadienyl radical)^{23,24} followed by loss of an anionic leaving group (Cl^-) to provide the α -heteroaryl amine product 5. This reduction step would simultaneously complete the photoredox cycle ($\text{Ir}(\text{II})$ to $\text{Ir}(\text{III})$) *via* reconstitution of the $\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ catalyst. Importantly, the incorporation of an anionic leaving group in the arene coupling partner would render this process redox neutral, obviating the need to employ either stoichiometric oxidants or reductants, while only generating HX as a reaction byproduct. Moreover, we anticipated that this strategy would enable the coupling of a diverse array of heteroaromatic rings while exhibiting complementary scope to our previously reported cyanoarene radical anion pathway.

Results and discussion

We initiated our α -amino arylation investigation by examining the utility of various 2-halo-benzothiazoles given their utility as potent electrophiles in two-electron $\text{S}_{\text{N}}\text{Ar}$ chemistry. While attempts to employ the cyano-substituted benzothiazole met with little success, we were delighted to find that bromo- and chloro-leaving groups provided the product of α -heteroarylation in useful yields using tris(2-phenylpyridinato- C^2, N)iridium(III)

Table 1 Evaluation of arene coupling partners and photocatalysts

Entry	Leaving group (X)	Amine (equiv.)	Photocatalyst (mol%)	Yield ^a (%)
1	CN	3	$\text{Ir}(\text{ppy})_3$ (1 mol%)	4%
2	I	3	$\text{Ir}(\text{ppy})_3$ (1 mol%)	5%
3	Br	3	$\text{Ir}(\text{ppy})_3$ (1 mol%)	50%
4	Cl	3	$\text{Ir}(\text{ppy})_3$ (1 mol%)	56%
5	F	3	$\text{Ir}(\text{ppy})_3$ (1 mol%)	16%
6	Cl	3	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (1 mol%)	75%
7	Cl	3	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (0.5 mol%)	78%
8	Cl	1.5	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (0.5 mol%)	77%
9 ^b	Cl	1.5	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (0.5 mol%)	84%
10 ^c	Cl	1.5	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (0.5 mol%)	87%
11	Cl	1.5	None	11%
12 ^d	Cl	1.5	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (0.5 mol%)	0%
13 ^e	Cl	1.5	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$ (0.5 mol%)	0%

^a Yield after 24 h determined by ¹H NMR analysis of crude reaction mixture with internal standard. Reactions performed with 2 equiv. NaOAc and 0.25 M DMA. ^b With 3 equiv. H₂O. ^c With 10 equiv. H₂O, isolated yield. ^d Performed in the absence of light. ^e Performed in the absence of NaOAc.

pyrimidine substrates were found to be viable, with a second electron-withdrawing substituent typically providing superior yields. Thus, 2-chloropyrimidines bearing ester, amide, or trifluoromethyl functionality at the 5-position undergo coupling with excellent efficiency (entries 1–3, 73–84% yield).²⁷ Pyrimidines that incorporate a second aromatic ring (entry 4, 73% yield) or a halide substituent (entry 5, 74% yield) may also be successfully employed. Performing these reactions at sub-ambient temperatures (0 °C) with added water is critical for suppressing arene decomposition and achieving high levels of reaction efficiency (see ESI†). Pyrimidines bearing a chloride leaving group at the 4-position also undergo coupling (entries 6–8, 72–92% yield), and *ortho*-substitution on the pyrimidine ring is tolerated, albeit with diminished yield (entry 9, 64% yield). Again, the compatibility of a second chloride substituent on the pyrimidine substrate allows for subsequent product functionalization *via* a range of transformations. Finally, this photoredox protocol may be used to install a variety of other 6-membered heteroarenes including purines (entry 10, 73% yield) and 1,3,5-triazines (entries 11 and 12, 88 and 84% yield).

With regard to the scope of the amine coupling partner, we anticipated that a range of readily oxidized tertiary anilines would serve as competent substrates. Indeed, using 2-chlorobenzoxazole as a representative heteroarene, we were pleased to find that both acyclic (Table 4, entries 1 and 2, 83 and 72% yield) and cyclic dialkylanilines (entries 3 and 7, 79 and 94% yield) undergo α -arylation with excellent efficiency. Moreover, saturated heterocyclic ring systems such as morpholine, thiomorpholine, and piperazine were found to be amenable to this photoredox heteroarylation (entries 4–6, 62–84% yield). Electron-donating as well as withdrawing substituents on the *N*-aryl ring are tolerated, and may be incorporated at the *para*, *meta*, or *ortho* positions (entries 8–12, 79–93% yield). Medicinally relevant tetrahydroquinoline and indoline architectures can also be successfully employed, with arylation proceeding regioselectively at the α -amino ring position *in lieu* of the exocyclic benzylic site (entries 13 and 14, 67 and 62% yield). Unsymmetrical, acyclic dialkylanilines interestingly undergo functionalization at methyl α -substituents in preference to methine (entry 15, single regioisomer) and methylene (entry 16, 9 : 1 r.r.) α -positions.²⁸ Finally, we provide examples of heteroaryl chloride-amine combinations that undergo coupling with moderate levels of reaction efficiency. As shown, the use of *N,N*-dimethylaniline, *N*-phenylmorpholine, and *N*-phenylpiperazine with 4,6-dichloropyrimidine resulted in diminished yields (entries 17–19, 22–66% yield). While these transformations remain of value to medicinal chemists, we are focused upon overcoming these limitations in efficiency.²⁹ To demonstrate the scalability of the arylation reaction, we additionally performed the coupling of 2-chlorobenzoxazole and *tert*-butyl-4-phenylpiperazine-1-carboxylate on a 4.0 mmol scale, providing the α -arylated product in 80% yield (1.22 g product, *cf.* Table 4, entry 6, 84% yield).

Mechanism

We have sought to obtain evidence for the proposed homolytic aromatic substitution mechanism as outlined in Scheme 1.

Table 4 Photoredox amine α -heteroarylation: amine scope^a

amine	heteroarene	α -heteroaryl amine
1: 83% yield		
(±)-2: 72% yield ^b		
(±)-3: 79% yield ^c		
(±)-4: 82% yield		
(±)-5: 62% yield ^d		
(±)-6: 84% yield		
(±)-7: 94% yield		
(±)-8: R = 4-OMe, 92% yield ^b		
(±)-9: R = 4-CF ₃ , 79% yield		
(±)-10: R = 4-Br, 88% yield ^b		
(±)-11: R = 3-Br, 86% yield		
(±)-12: R = 2-F, 93% yield ^b		
(±)-13: 67% yield ^c		
(±)-14: 62% yield		
15: 75% yield		
16: 84% yield, 9:1 r.r.		
17: 66% yield ^{e,f}		
(±)-18: 22% yield ^{e,g}		
(±)-19: 35% yield ^{e,s}		

^a Conditions employ 10 equiv. H₂O unless otherwise noted. ^b Performed with 5 equiv. H₂O. ^c Performed at 0 °C. ^d Performed with 15 equiv. H₂O. ^e Performed with 1.0 equiv. amine and 1.5 equiv. arene at 0 °C. ^f Performed with 40 equiv. H₂O. ^g Performed with 20 equiv. H₂O.

Stern–Volmer fluorescence quenching studies using the preferred Ir(ppy)₂(dtbbpy)PF₆ photocatalyst revealed significant emission quenching by *N*-phenylpyrrolidine, yet no change in the emission was observed using a variety of chloroheteroarene substrates. This finding indicates that a reductive quenching

cycle is operative (wherein the excited state of the catalyst first performs amine oxidation). To determine if the neutral α -amino radical **1** undergoes addition to the ground state heteroarene **3** (as proposed in Scheme 1) or if the chloroarene undergoes reduction *via* electron transfer with the transient Ir^{III}(ppy)₂(dtbbpy) species (as observed in our cyanoarene studies),¹⁴ the reduction potentials of several chloroheteroarene substrates were measured by cyclic voltammetry. These potentials were found to range from -1.70 V to -2.19 V *vs.* SCE (see ESI†). More specifically, 2-chlorobenzoxazole was found to undergo reduction at -2.19 V *vs.* SCE, indicating that electron transfer from the Ir^{III}(ppy)₂(dtbbpy) species is highly endothermic and suggesting that a radical anion–neutral radical coupling mechanism is not operative.^{30,31}

Conclusions

In conclusion, we have leveraged photoredox catalysis to achieve the α -heteroarylation of tertiary amines with a diverse array of five- and six-membered heterocycles *via* a putative homolytic aromatic substitution pathway. Given the prevalence of heterocycles in medicinal agents, as well as the commercial availability of a large number of chloroheteroarenes, we expect that this reaction will find immediate application in the synthesis of biologically active compounds.

Acknowledgements

Financial support was provided by the NIHGMS (R01 01 GM093213-01), Merck and Amgen. C. K. P. is grateful for a Bristol-Myers Squibb graduate fellowship. We thank Yuan Hu for generous assistance with cyclic voltammetry.

Notes and references

- 1 T. J. Ritchie, S. J. F. Macdonald, S. Peace, S. D. Pickett and C. N. Luscombe, *MedChemComm*, 2012, **3**, 1062.
- 2 N. A. McGrath, M. Brichtacek and J. T. Njardarson, *J. Chem. Educ.*, 2010, **87**, 1348.
- 3 (a) M. B. Smith and J. March, *Advanced Organic Chemistry*, Wiley-Interscience, New York, 5th edn, 2001, p. 850; (b) J. A. Joule and K. Mills, *Heterocyclic Chemistry*, Wiley, Chichester, 5th edn, 2010; (c) J. F. Bunnett and R. E. Zahler, *Chem. Rev.*, 1951, **49**, 273; (d) J. F. Bunnett, *J. Chem. Educ.*, 1974, **51**, 312; (e) K. Walsh, H. F. Sneddon and C. J. Moody, *ChemSusChem*, 2013, **6**, 1455.
- 4 (a) J. P. Wolfe, S. Wagaw, J.-F. Marcoux and S. L. Buchwald, *Acc. Chem. Res.*, 1998, **31**, 805; (b) D. S. Surry and S. L. Buchwald, *Chem. Sci.*, 2011, **2**, 27; (c) J. F. Hartwig, *Angew. Chem., Int. Ed.*, 1998, **37**, 2046; (d) G. Evans, N. Blanchard and M. Toumi, *Chem. Rev.*, 2008, **108**, 3054.
- 5 (a) S. D. Roughley and A. M. Jordan, *J. Med. Chem.*, 2011, **54**, 3451; (b) J. S. Carey, D. Laffan, C. Thomson and M. T. Williams, *Org. Biomol. Chem.*, 2006, **4**, 2337.
- 6 C. K. Donawho, Y. Luo, Y. Luo, T. D. Penning, J. L. Bauch, J. J. Bouska, V. D. Bontcheva-Diaz, B. F. Cox, T. L. DeWeese, L. E. Dillehay, D. C. Ferguson, N. S. Ghoreishi-Haack, D. R. Grimm, R. Guan, E. K. Han, R. R. Holley-Shanks, B. Hristov, K. B. Idler, K. Jarvis, E. F. Johnson, L. R. Kleinberg, V. Klinghofer, L. M. Lasko, X. Liu, K. C. Marsh, T. P. McGonigal, J. A. Meulbroeck, A. M. Olson, J. P. Palma, L. E. Rodriguez, Y. Shi, J. A. Stavropoulos, A. C. Tsurutani, G.-D. Zhu, S. H. Rosenberg, V. L. Giranda and D. J. Frost, *Clin. Cancer Res.*, 2007, **13**, 2728.
- 7 N. H. Huel, H. Nar, H. Priepeke, U. Ries, J.-M. Stassen and W. Wienen, *J. Med. Chem.*, 2002, **45**, 1757.
- 8 M. Reuveni, *Eur. J. Plant Pathol.*, 2003, **109**, 243.
- 9 (a) R. K. Dieter and S. Li, *Tetrahedron Lett.*, 1995, **36**, 3613; (b) R. K. Dieter and S. Li, *J. Org. Chem.*, 1997, **62**, 7726.
- 10 (a) K. R. Campos, A. Klapars, J. H. Waldman, P. G. Dormer and C. Chen, *J. Am. Chem. Soc.*, 2006, **128**, 3538; (b) A. Klapars, K. R. Campos, J. H. Waldman, D. Zewge, P. G. Dormer and C. Chen, *J. Org. Chem.*, 2008, **73**, 4986; (c) G. Barker, P. O'Brien and K. R. Campos, *Org. Lett.*, 2010, **12**, 4176; (d) G. Barker, J. L. McGrath, A. Klapars, D. Stead, G. Zhou, K. R. Campos and P. O'Brien, *J. Org. Chem.*, 2011, **76**, 5936.
- 11 N. Yoshikai, A. Mieczkowski, A. Matsumoto, L. Ilies and E. Nakamura, *J. Am. Chem. Soc.*, 2010, **132**, 5568.
- 12 For reviews on amine α -functionalization: (a) K. R. Campos, *Chem. Soc. Rev.*, 2007, **36**, 1069; (b) E. A. Mitchell, A. Peschiulli, N. Lefevre, L. Meerpoel and B. U. W. Maes, *Chem.–Eur. J.*, 2012, **18**, 10092.
- 13 (a) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (b) J. W. Tucker and C. R. J. Stephenson, *J. Org. Chem.*, 2012, **77**, 1617; (c) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102; (d) F. Teplý, *Collect. Czech. Chem. Commun.*, 2011, **76**, 859.
- 14 (a) A. McNally, C. K. Prier and D. W. C. MacMillan, *Science*, 2011, **334**, 1114; (b) A subsequent variant was reported using trialkylamines and aryl halides: A. Singh, A. Arora and J. D. Weaver, *Org. Lett.*, 2013, **15**, 5390.
- 15 P. Renaud and L. Giraud, *Synthesis*, 1996, 913.
- 16 (a) F. Minisci, *Synthesis*, 1973, 1; (b) F. Minisci, *Top. Curr. Chem.*, 1976, **62**, 1; (c) F. Minisci, E. Vismara and F. Fontana, *Heterocycles*, 1989, **28**, 489; (d) F. Minisci, F. Fontana and E. Vismara, *J. Heterocycl. Chem.*, 1990, **27**, 79; (e) M. A. J. Dunton, *MedChemComm*, 2011, **2**, 1135.
- 17 (a) S. E. Vaillard, B. Schulte and A. Studer, *Radical-Based Arylation Methods in Modern Arylation Methods*, ed. L. Ackermann, Wiley-VCH, Weinheim, 2009, pp. 475–511; (b) A. Studer and M. Bossart, *Homolytic Aromatic Substitutions in Radicals in Organic Synthesis*, ed. P. Renaud and M. P. Sibi, Wiley-VCH, Weinheim, 2001, vol. 2, pp. 62–80; (c) W. R. Bowman and J. M. D. Storey, *Chem. Soc. Rev.*, 2007, **36**, 1803; (d) J. G. Traynham, *Chem. Rev.*, 1979, **79**, 323.
- 18 For examples of homolytic aromatic substitution reactions employing photoredox catalysis: (a) H. Cano-Yelo and A. Deronzier, *J. Chem. Soc., Perkin Trans. 2*, 1984, 1093; (b) L. Furst, B. S. Matsuura, J. M. R. Narayanam, J. W. Tucker and C. R. J. Stephenson, *Org. Lett.*, 2010, **12**, 3104; (c) D. A. Nagib and D. W. C. MacMillan, *Nature*, 2011, **480**,

- 224; (d) D. P. Hari, P. Schroll and B. König, *J. Am. Chem. Soc.*, 2012, **134**, 2958.
- 19 For other relevant examples of homolytic aromatic substitution: (a) M. Fiorentino, L. Testaferrri, M. Tiecco and L. Troisi, *J. Chem. Soc., Chem. Commun.*, 1977, 316; (b) C. J. Cowden, *Org. Lett.*, 2003, **5**, 4497; (c) Y. Fujiwara, J. A. Dixon, F. O'Hara, E. D. Funder, D. D. Dixon, R. A. Rodriguez, R. D. Baxter, B. Herlé, N. Sach, M. R. Collins, Y. Ishihara and P. S. Baran, *Nature*, 2012, **492**, 95.
- 20 (a) J. D. Slinker, A. A. Gorodetsky, M. S. Lowry, J. Wang, S. Parker, R. Rohl, S. Bernhard and G. G. Malliaras, *J. Am. Chem. Soc.*, 2004, **126**, 2763; (b) M. S. Lowry, J. I. Goldsmith, J. D. Slinker, R. Rohl, R. A. Pascal, Jr, G. G. Malliaras and S. Bernhard, *Chem. Mater.*, 2005, **17**, 5712.
- 21 E. T. Seo, R. F. Nelson, J. M. Fritsch, L. S. Marcoux, D. W. Leedy and R. N. Adams, *J. Am. Chem. Soc.*, 1966, **88**, 3498.
- 22 A pK_a of 18.8 has been calculated for the radical cation of *N,N*-dimethylaniline in acetonitrile: G. W. Dombrowski, J. P. Dinnocenzo, P. A. Zielinski, S. Farid, Z. M. Wosinska and I. R. Gould, *J. Org. Chem.*, 2005, **70**, 3791, The acetate base employed in this work may perform this deprotonation in organic solvents; the pK_a of acetic acid in acetonitrile has been determined to be 23.5: A. Kütt, I. Leito, I. Kaljurand, L. Sooväli, V. M. Vlasov, L. M. Yagupolskii and I. A. Koppel, *J. Org. Chem.*, 2006, **71**, 2829.
- 23 (a) J. Lilie, G. Beck and A. Henglein, *Ber. Bunsenges. Physik. Chem.*, 1971, **75**, 458; (b) M. Grätzel, A. Henglein, J. Lilie and M. Scheffler, *Ber. Bunsenges. Physik. Chem.*, 1972, **76**, 67.
- 24 While a cyclohexadienyl radical is not a perfect approximation for the heterocyclic dienyl radical **4**, we anticipate that the heterocyclic radical will undergo reduction even more readily.
- 25 For more data on the effect of water, see ESI.† For another example of the beneficial role of water in single-electron transfer chemistry, see: T. Hoshikawa and M. Inoue, *Chem. Sci.*, 2013, **4**, 3118.
- 26 Reactivity in the absence of photocatalyst may involve electron transfer from excited electron donor–acceptor complexes, see: (a) R. Foster, *J. Phys. Chem.*, 1980, **84**, 2135; (b) E. Arceo, I. D. Jurberg, A. Álvarez-Fernández and P. Melchiorre, *Nat. Chem.*, 2013, **5**, 750.
- 27 While we have previously demonstrated the installation of pyridine rings using the corresponding cyanoarenes (ref. 14), our efforts to couple more electron-deficient heteroaryl nitriles (such as cyanopyrimidines) have thus far been unsuccessful.
- 28 See ESI† for a proposed model for the observed regioselectivity.
- 29 We postulate that the superior reactivity of pyrrolidines arises from the ability of this ring system to accommodate conjugation between the α -amino SOMO and the nitrogen lone pair without incurring destabilizing eclipsing interactions. See: (a) D. D. M. Wayner, K. B. Clark, A. Rauk, D. Yu and D. A. Armstrong, *J. Am. Chem. Soc.*, 1997, **119**, 8925; (b) D. Griller, J. A. Howard, P. R. Marriott and J. C. Scaiano, *J. Am. Chem. Soc.*, 1981, **103**, 619.
- 30 Stern–Volmer quenching experiments were also performed using *fac*-Ir(ppy)₃, a complex which in its photoexcited state ($E_{1/2}^{IV*/III} = -1.73$ V vs. SCE) is slightly more reducing than Ir^{III}(ppy)₂(dtbbpy) ($E_{1/2}^{III/II} = -1.51$ V vs. SCE). Minimal quenching ($k_q\tau_0 < 2$ M⁻¹) was observed in the presence of pyrimidine and purine substrates, and no quenching was detected in the presence of 2-chlorobenzoxazole (see ESI†).
- 31 It has furthermore been demonstrated that the reduction of chloroheteroarenes to the corresponding radical anions typically results in halide fragmentation to generate aryl radicals. For examples of this reactivity, see: D. R. Carver, A. P. Komin, J. S. Hubbard and J. F. Wolfe, *J. Org. Chem.*, 1981, **46**, 294.